



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

A prospective, phase II, single-centre, cross-sectional, randomised study investigating Dehydroepiandrosterone supplementation and its Profile in Trauma: ADaPT

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2020-040823
Article Type:	Protocol
Date Submitted by the Author:	03-Jun-2020
Complete List of Authors:	<p>Bentley, Conor; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences</p> <p>Potter, Claire; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Yakoub, Kamal; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Brock, Kristian; University of Birmingham College of Medical and Dental Sciences, D3B</p> <p>Toman, Emma; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction and Microbiology Research Centre</p> <p>Taylor, Angela; University of Birmingham College of Medical and Dental Sciences, Institute of Metabolism and Systems Research</p> <p>Shaheen, Fozia; University of Birmingham College of Medical and Dental Sciences, Institute of Metabolism and Systems Research</p> <p>Gilligan, Lorna; University of Birmingham College of Medical and Dental Sciences, D3B</p> <p>Athwal, Amrita; NIHR Birmingham Liver Biomedical Research Unit Clinical Trials Group (D3B team), CRUK Clinical Trials Unit, University of Birmingham, Birmingham, UK</p> <p>Barton, Darren; NIHR Birmingham Liver Biomedical Research Unit Clinical Trials Group (D3B team), CRUK Clinical Trials Unit, University of Birmingham, Birmingham, UK</p> <p>Carrera, Ronald; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Young, Katie; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Desai, Amisha; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>McGee, Kirsty; University of Birmingham College of Medical and Dental Sciences, Institute of Metabolism and Systems Research</p> <p>Ermogenous, Christos; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Greig, Carolyn; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences; MRC-Arthritis Research UK Centre for Musculoskeletal Ageing Research</p> <p>Hazeldine, Jon; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Arlt, Wiebke; University of Birmingham College of Medical and Dental</p>

	Sciences, Institute of Metabolism and Systems Research; National Institute of Health Research Birmingham Biomedical Research Unit Lord, Janet; University of Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre; National Institute of Health Research Birmingham Biomedical Research Unit Foster, Mark; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre; Royal Centre for Defence Medicine
Keywords:	TRAUMA MANAGEMENT, Adult intensive & critical care < INTENSIVE & CRITICAL CARE, Sex steroids & HRT < DIABETES & ENDOCRINOLOGY

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A prospective, phase II, single-centre, cross-sectional, randomised study investigating Dehydroepiandrosterone supplementation and its Profile in Trauma: ADaPT

Conor Bentley ^{1,2,3}, Claire Potter ⁴, Kamal M. Yakoub ¹ Kristian Brock ⁴, Emma Toman ¹, Angela E. Taylor ⁵, Fozia Shaheen ⁵, Lorna C. Gilligan ⁵, Amrita Athwal ⁴, Darren Barton ⁴, Ronald Carrera ¹, Katie Young ¹, Amisha Desai ¹, Kirsty McGee ^{1,3}, Christos Ermogenous ^{1,3}, Carolyn Greig ^{2,6}, Jon Hazeldine ^{1,3}, Wiebke Arlt ^{5,6}, Janet M. Lord ^{1,3,6}, Mark A. Foster ^{1,3,7}

¹ NIHR Surgical Reconstruction and Microbiology Research Centre, University Hospital Birmingham, Queen Elizabeth Hospital Birmingham, Birmingham B15 2WB, UK.

² School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, UK.

³ MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research, Institute of Inflammation and Ageing, Birmingham University Medical School, Birmingham B15 2TT, UK.

⁴ Diagnostics, Drugs, Devices and Biomarkers team (D³B) Cancer Research Clinical Trials Unit, University of Birmingham, Birmingham, UK.

⁵ Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK.

⁶ NIHR Biomedical Research Centre, University Hospital Birmingham and University of Birmingham, Birmingham, UK.

⁷ Royal Centre for Defence Medicine, Birmingham Research Park, Birmingham B15 2SQ, UK.

1
2
3 22
4
5
6
7 23
8
9 24
10
11
12 25
13
14
15 26
16
17
18
19 27
20
21 28
22
23
24 29
25
26 30
27
28 31
29
30 32
31
32 33
33
34 34
35
36 35
37
38 36
39
40 37
41
42 38
43
44 39
45
46 40
47
48 41
49
50
51
52
53
54
55
56
57
58
59
60

Correspondence to: Conor Bentley, NIHR Surgical Reconstruction and Microbiology Research

Centre, University Hospital Birmingham, Birmingham B15 2TH, UK.

email: conor.bentley@nhs.net

Tel: 0121 371 4242

Word Count: 4146

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Introduction: The improvements in short-term outcome after severe trauma achieved through early resuscitation and acute care can be offset over the following weeks by an acute systemic inflammatory response with immuneparesis leading to infection, multi-organ dysfunction/failure (MOD/MOF) and death. Serum levels of the androgen precursor dehydroepiandrosterone (DHEA) and its sulphated ester DHEAS, steroids with immune-enhancing activity, are low after traumatic injury at a time when patients are catabolic and immunosuppressed. Addressing this deficit and restoring the DHEA(S) ratio to cortisol may provide a range of physiological benefits, including immune modulatory effects.

Objective: Our primary objective is to establish a dose suitable for DHEA supplementation in patients after acute trauma to raise circulating DHEA levels to at least 15 nmol/L. Secondary objectives are to assess if DHEA supplementation has any effect on neutrophil function, metabolic and cytokine profiles and which route of administration (oral vs sub-lingual) is more effective in restoring circulating levels of DHEA, DHEAS and downstream androgens.

Methods and analysis: ADaPT is a prospective, phase II, single-centre, cross-sectional, randomised study with a planned recruitment between between April 2019 and July 2021 that will investigate DHEA supplementation and its effect on serum DHEA, DHEAS and downstream androgens in trauma. A maximum of 270 patients will receive sublingual or oral DHEA at 50, 100 or 200 mg daily over 3 days. Females aged ≥ 50 years with neck of femur fracture and male and female major trauma patients, aged 16-50 years with an injury severity score ≥ 16 , will be recruited.

Ethics and dissemination: This protocol has been approved by a Research Ethics Committee (Reference 18/WM/0102) on 8th June 2018. Results will be disseminated via peer-reviewed publications and presented at national and international conferences.

66

Trial registration: This trial is registered with the European Medicines Agency (EudraCT: 2016-004250-15) and ISRCTN (12961998). It has also been adopted on the National Institute of Health Research portfolio (CPMS ID:38158).

Trial Progression: The study recruited its first patient on 2nd April 2019 and held its first data monitoring committee on 8th November 2019. As of May 2020 there were 23 enrolled patients, with both male cohorts increasing to 100 mg. All female groups remain on 50 mg DHEA.

74

Article Summary

Strengths and limitations of this study

►► A phase II experimental study of a food supplement in the USA, but a class C controlled drug in the EU, in an acutely injured cohort with a large unmet need for new and effective therapies in both the short and long-term recovery from injury.

►► Aims to address what dose and route of administration are required to normalise DHEA levels in a cohort that we know have low levels, and any added benefit this restoration has upon immune and inflammatory parameters.

►► The rapid turnaround time from bedside to bench and from bench to interim analysis will minimise inappropriate dosing and expenditure of public money on ineffective doses. Thus, limiting patient exposure to insufficient dosing and unnecessary specimen collection.

Keywords: steroids, geriatric trauma, major trauma, endocrine, DHEA; DHEAS; immune function

88

89 Introduction

90 The improvements in short-term outcome after severe trauma achieved through early
91 resuscitation and acute care are offset over the following weeks by an acute systemic
92 inflammatory response with immuneparesis leading to infection, multi-organ
93 dysfunction/failure (MOF) and death^{1,2}. The combination of excessive pro- and anti-
94 inflammatory responses impair the rehabilitation phase of trauma, including wound healing,
95 physical and emotional recovery³. Upregulation of glucocorticoid biosynthesis promotes a
96 catabolic state, lasting several weeks and associated with a significant reduction in muscle
97 mass⁴.

98 Analysis of gender differences in trauma has shown that women, particularly those under 30
99 years of age, have fewer infections and a better outcome from sepsis than men^{5,6}. The
100 protective effects of oestrogens on immune cells and organ function highlight the potential
101 role of sex hormones in modulating trauma related immune dysfunction⁷. Cellular immunity
102 is also influenced by hormone production, and members of our group have shown that the
103 adrenal response to stress, specifically the ratio of cortisol to dehydroepiandrosterone
104 sulphate (DHEAS), is linked to neutrophil bactericidal function, specifically superoxide
105 production^{8,9}.

106 Dehydroepiandrosterone (DHEA) and its sulphate ester DHEAS are the most abundant
107 steroids in the human circulation; DHEA is a precursor of sex hormones modulating several
108 physiologic processes including metabolism, muscle protein synthesis and cardiovascular
109 function^{10,11}. DHEA is converted to active androgens in peripheral target cells including
110 immune cells¹² and is also converted to DHEAS by the action of DHEA sulfotransferase
111 (SULT2A1)¹³. We have shown that DHEA sulphation is down-regulated in acute inflammation
112 systemic inflammatory response syndrome (SIRS) and sepsis¹⁴. Hazeldine et al reviewed the

113 numerous immune effects of DHEA ¹⁵ highlighting how DHEAS, but not DHEA, enhances
114 neutrophil superoxide generation via a protein kinase C (PKC) mediated pathway, a vital
115 immune response in fighting bacterial infections¹⁶.

116 The roles of DHEA and DHEAS in severe injury are relatively unexplored. The majority of
117 studies focus upon cortisol responses¹⁷, whereas our data suggest that it is the cortisol: DHEAS
118 ratio post-trauma that has a superior prognostic ability^{18,19}. Although critical for survival,
119 prolonged hypercortisolaemia is known to be detrimental in part due to its
120 immunosuppressive effects²⁰. This intra-adrenal shift causes decreased levels of circulating
121 testosterone and oestrogen, resulting in rapid lean body mass loss²¹, in addition to increased
122 susceptibility to infection and sepsis²². Correcting the cortisol: DHEA or cortisol: DHEAS ratio
123 via the administration of DHEA has yet to be undertaken in traumatically-injured patients
124 despite DHEA and DHEAS being below the reference ranges for 6 weeks and 6 months post-
125 injury respectively⁴.

126 DHEA is subject to first-pass metabolism in healthy individuals, which in turn may lead to a
127 non-physiological metabolism after an oral dose²³. Bypassing first-pass metabolism using a
128 sublingual or buccal preparation²⁴ should improve the bioavailability of DHEA ^{25,26}. Previous
129 studies in the healthy older people have shown that supplementation with 50 mg DHEA orally
130 once daily can restore both serum DHEA and DHEAS levels to that of men and women in the
131 third decade of life^{23,27}. Moreover, the current literature suggests that DHEA supplementation
132 may not only be beneficial for immune function but extend to bone remodelling, muscle
133 composition, psychological and neurological improvements²⁸.

134 This study will seek to estimate the optimal dose and route of administration in trauma
135 patients of DHEA to increase serum levels of DHEA and DHEAS to those seen in healthy adults.

136 We also aim to find the optimal dose to enhance immune function, as demonstrated through

1
2
3 137 changes in neutrophil phagocytosis and ROS production. The pilot data generated from the
4
5
6 138 study is necessary to develop the protocol for a randomised controlled trial that will
7
8 139 determine whether DHEA supplementation may improve outcomes in injured patients.
9

10
11 140
12
13 141
14

15
16 142 **Rationale**
17

18
19 143 **Justification of the patient population**

20 144 Although improvements in short-term outcomes from traumatic injury via aggressive early
21
22
23 145 resuscitation and acute care, over 5 million people worldwide die each year from serious
24
25 146 injury²⁹. With improved pre-hospital medicine mitigating the immediate threats to life, it is
26
27
28 147 the following weeks after traumatic injury, that has seen the dysregulated SIRS in susceptible
29
30 148 patients. The SIRS response is accompanied by a compensatory anti-inflammatory response
31
32
33 149 (CARS)³⁰ to restore homeostasis. SIRS and CARS may progress to the persistent inflammation,
34
35 150 immunosuppression and catabolism syndrome (PICS)^{31,32}. PICS further compounds the insult
36
37
38 151 from the initial injury and results in an increased risk of infection, MOF and late deaths^{1,2}.
39
40 152 Therefore, strategies to mitigate these detrimental outcomes for patients are needed in the
41
42
43 153 short, medium and long-term care of trauma patients.

44
45 154 The young trauma cohort will be recruited alongside a cohort of older (≥ 50 years old) female
46
47
48 155 patients who have sustained a hip fracture at the neck of femur (NOF). According to the
49
50 156 National Hip Fracture Database (2018), the NOF burden upon the UK economy is estimated
51
52 157 to be around £1bn per annum³³. This considerable cost is set to rise as the population ages.
53
54
55 158 By the age of 40, decreasing serum levels of DHEA and DHEAS are observed in both sexes^{34,35};
56
57
58 159 a phenomenon sometimes referred to as “adrenopause”. Circulating levels of DHEA and
59
60 160 DHEAS are lower in women than in men; however, in post-menopausal women, adrenal

androgens are a source of almost all active oestrogens³⁶. To the best of our knowledge, there has been no traumatic injury or NOF interventional studies supplementing DHEA or DHEAS, despite its therapeutic potential in the immediate to longer-term care of the young and aged trauma patient. In this pilot study, we will recruit a female NOF cohort as well as a young adult trauma cohort with both male and female patients presenting at the MTC and Intensive Care Unit (ICU) at QEHB.

Choice of treatment

The DHEA doses chosen in this study (50 mg, 100 mg and 200 mg) were selected based upon *in vivo* studies that demonstrated these doses to be both safe and effective at raising the levels of DHEA to that of healthy young adult levels³⁷. DHEA is at its highest in the third decade of life. After which there is a steady decrease over the life course in both males and females. These doses have also previously been used in adrenal insufficiency³⁸, older people³⁹, and young males⁴⁰ and females²⁴, causing transient hyperandrogenism with acne being the most frequently reported side effect^{41,42}. DHEA is activated to downstream androgens by stepwise conversion catalysed by several steroidogenic enzymes. However, we do not know whether the expression and activity of these enzymes are affected by major trauma and inflammation. There is evidence that inflammation and trauma affect DHEA sulfation^{4,14} and may shift the conversion of DHEA to a higher rate of androgen activation.

Trauma and inflammation can impact adversely on gut absorption⁴³ and hepatic first-pass metabolism⁴⁴. Therefore, we have chosen to administer DHEA as a sublingual preparation. Sub-lingual administration is commonly used by nursing staff in the ICU context, especially in those patients who have contraindications to oral administration or by patients wishing to self-administer⁴⁵.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

OBJECTIVES

Primary Objective

The primary objective of this study is to establish the daily dose of DHEA that restores serum DHEA levels to at least 15 nmol/L, i.e. the mid healthy adult reference range, in trauma and hip fracture patients after 3 days of supplementation.

Secondary Objective

Secondary objectives include observing the effect of single and multiple DHEA doses on circulating DHEA, DHEAS and downstream androgens. Additionally, we will investigate whether route of DHEA administration (oral vs sublingual) modulates the profile of circulating DHEA. This will be determined by assessing the efficacy of each route via statistical modelling. The immune response to the DHEA supplementation will be assessed as a marker of potential clinical relevance.

METHODS

Trial Design

ADaPT is a prospective, randomised, open-label, trial conducted in male and female adult trauma patients and older females who have suffered a fracture of the neck of the femur. This is a single-site study with patients recruited from Queen Elizabeth Hospital, Birmingham, UK (QEHB). Three doses of DHEA will be investigated in this trial: 50, 100, 200 mg, each administered once daily for 3 days via either oral or sublingual tablets. The trial has an adaptive design in order to answer both the primary and secondary objectives, with regular interim analysis to minimise the investigation of inactive doses. The trial consists of two components. The first component of the trial is to determine the dose of DHEA needed to sufficiently raise serum DHEA levels to at least 15 nmol/L after 3 days of DHEA administration.

209 Based on previous work, 15 nmol/L has been selected as this is the lower acceptable level of
210 DHEA observed in healthy young adults. Our recent analysis of the steroid response to trauma
211 has shown that DHEA levels were very low and often undetectable for several weeks after
212 trauma ⁴. The second component of the trial is to investigate if DHEA will enhance neutrophil
213 function. The study was approved by The Medicines and Healthcare products Regulatory
214 Agency (MHRA) for the use of DHEA as an investigational medicinal product. Ethical approval
215 was obtained from the West Midlands Research Ethics Committee (Reference 18/WM/0102).
216 The trial will be conducted in accordance with the Declaration of Helsinki (World Medical
217 Association 2008). **Figure 1** summarises the patient pathway of the trial. The protocol (version
218 5.0, 28th June 2019) has been prepared in accordance with the SPIRIT guidelines ⁴⁶.

221 Patient selection

222 A maximum of 270 patients will be enrolled across three patient groups (young male trauma,
223 young female trauma and female hip fracture). These trauma patient groups have been
224 selected due to the immuneparesis effects caused by the acute systemic inflammatory
225 response that follows severe trauma. The hip fracture group was selected as they have low
226 serum DHEA and DHEAS due to adrenopause, and there are several papers showing an
227 association between HPA axis activity measures and outcomes in these patients^{9,19,47,48}.
228 Patients admitted to QEHB will all be pre-screened and assessed for eligibility. Patients will
229 be screened between 07:00 – 20:00, 7 days a week. Potential participants will have an
230 assessment of their medical history and current trauma injury and the eligible patients will be
231 recruited. The eligibility criteria have been split into trauma patients and hip fracture patients
232 (**Table 1**). The study will not exclude NOF patients with dementia where supplementation is

233 currently being tested in the prevention and treatment of age-related cognitive impairment
234 without deleterious effects⁴⁹.

Table 1 Patient inclusion and exclusion criteria

Trauma patients inclusion criteria	
-	Aged 16 - 50 years of age
-	Severely injured trauma patient (Injury severity score (ISS) ≥16 and ≤50)
-	Admitted to University Hospital Birmingham within 6 days of trauma
-	Anticipated to be an inpatient for the 12-day trial period
Trauma patients exclusion criteria	
-	Ages <16 or >51
-	ISS <16 or >50
-	Isolated brain injury
-	Unlikely to survive the study period
-	Known hormone sensitive malignancy
-	Known Prostatic hypertrophy (M)
-	Female patients taking HRT medication
-	Intake of any drugs that interfere with adrenal function in the last 3 months:
Increased metabolism of glucocorticoids	
•	corticosteroids
Impaired glucocorticoid action	
Peripheral glucocorticoid insensitivity	
•	Glucocorticoid receptor antagonist—mifepristone.
•	Suppression of glucocorticoid-induced gene transcription—chlorpromazine, imipramine.
Inhibition of steroidogenic enzymes involved in cortisol production	
•	Inhibition of mitochondrial (type 1) cytochrome P450 enzymes (CYP11A1, CYP11B1, CYP11B12)—ketoconazole, fluconazole, itraconazole, etomidate, metyrapone, aminoglutethimide.
•	Inhibition of 3β-HSD2—trilostane.
-	Pre-existing liver impairment or chronic liver failure
-	Previous or current malignancy or invasive cancer diagnosed within the past 3 years except for adequately treated basal cell and squamous cell carcinoma of the skin and in situ carcinoma of the uterine cervix
-	
-	Pregnant and/or breast-feeding females (women of childbearing potential to complete serum pregnancy test)
-	Known hypersensitivity to the active substance or excipient
-	Known thromboembolic events in the last 12 months and any predisposition to thrombosis e.g. factor V leiden
Hip fracture patients inclusion criteria	
-	Aged 50 years and older
-	Female
-	Neck of femur fracture

-	Admitted to University Hospital Birmingham within 6 days of fracture
-	Anticipated to be an inpatient for the 12 day trial period
Hip Fracture patients exclusion criteria	
-	<50 years old
-	Unlikely to survive the study period
-	Previous or known hormone sensitive malignancy
-	Intake of any drugs that interfere with adrenal function in the last 3 months:
Increased metabolism of glucocorticoids	
•	Concomitant use reduces corticosteroid concentrations
Impaired glucocorticoid action	
Peripheral glucocorticoid insensitivity	
•	Glucocorticoid receptor antagonist—mifepristone.
•	Suppression of glucocorticoid-induced gene transcription—chlorpromazine, imipramine.
Inhibition of steroidogenic enzymes involved in cortisol production	
•	Inhibition of mitochondrial (type 1) cytochrome P450 enzymes (CYP11A1, CYP11B1, CYP11B12)—ketoconazole, fluconazole, itraconazole, etomidate, metyrapone, aminoglutethimide.
•	Inhibition of 3 β -HSD2—trilostane.
-	
-	Pre-existing liver impairment or chronic liver failure
-	Previous or current malignancy or invasive cancer diagnosed within the past 3 years except for adequately treated basal cell and squamous cell carcinoma of the skin and in situ carcinoma of the uterine cervix
-	Pregnant and/or breastfeeding (women of childbearing potential to complete serum pregnancy test)
-	Known hypersensitivity to the active substance or excipient
-	Females on Hormone Replacement Therapy medication
-	Known thromboembolic events in the last 12 months and any predisposition to thrombosis e.g. factor V leiden

Randomisation

Patients who meet the eligibility criteria and provide consent are randomised to receive DHEA via either oral tablets or sublingual tablets once daily for 3 days using a 1:1 randomisation ratio. With three populations of patients (male-trauma, female-trauma and female-hip fracture) and two routes of administration, there will be 6 groups in total. Randomisation will take place using a secure web-based tool. Nursing and medical staff will use the Clinical RESearch Tool (CREST), developed at University Hospitals Birmingham Foundation Trust

1
2
3 243 (UHBFT), to randomly assign patients and provide an anonymised electronic case report form,
4
5
6 244 for trial management, data collection and adverse event reporting. The prevailing dose of
7
8 245 DHEA (50, 100 or 200 mg) for administration in a group will be adapted in response to
9
10 246 sequential analysis of interim outcomes. Each group will have its own dose. Blinding is not
11
12
13 247 possible as the difference in DHEA delivery method is evident to both the clinician and the
14
15 248 participant. If a contraindication to oral or the sublingual route present prior to commencing
16
17
18 249 the study intervention, forced randomisation will occur.
19
20
21 250
22
23
24

25 251 **Study Intervention**

26 252 The supplementation of DHEA will commence in the second week after injury, which has
27
28 253 previously been shown to be a time when trauma patients become maximally unwell as a
29
30 254 result of sepsis and multiple organ failure⁵⁰. Additionally, this time point has been selected to
31
32
33 255 optimise patient recruitment from both the NOF cohort and trauma patients (median stay 18
34
35 256 days s 13 days respectively) both DHEA and DHEAS levels post-injury ⁵¹. Recruiting an in-
36
37
38 257 hospital cohort provides an opportunity to monitor patients and assess the impact that this
39
40 258 class C controlled drug has upon the serial steroid profile and immune function during a
41
42
43 259 period of vulnerability over 3 days of administration.
44
45
46 260
47
48

49 261 **Participant timeline**

50 262 The trial intervention will occur over three days, during the first 12 days while inpatients at
51
52
53 263 QEHB. Three doses of DHEA will be investigated in all patient, and treatment will occur on day
54
55 264 8, 9 and 10 only. All cohorts will begin the study on 50 mg, and the dose administered to the
56
57
58 265 next eligible patient to be included within a cohort will be escalated when interim analysis
59
60

determines if the primary objective has been reached. At no point will any patient escalate once they have been allocated a dose of DHEA.

Dose escalation

Dose escalation for the DHEA restoration part of the trial is dependent upon the serum DHEA levels. A dose that restores serum DHEA to ≥ 15 nmol/L (referred to as 'normalise DHEA') is sought in at least 13-out-of-15 patients or approximately 85% of patients. The decision to escalate dose in a cohort will be driven in the main by the outcomes of the patients in that cohort. However, valuable additional information will be garnered from the outcomes of patients in other cohorts. Flexible information sharing across related groups will be achieved using hierarchical regression models.

Once a dose has been established to normalise DHEA levels within a group and at least $n = 15$ have completed it, the DHEA will be escalated to the next dose to satisfy the second component of the trial; to determine whether further increases in DHEA supplementation will enhance immune function. The immune response component will focus on neutrophil phagocytosis and ROS production which will involve fewer research samples. $N = 15$ patients (within a cohort) will complete the immune response part of the trial at each subsequent dose escalation. If 50 mg is established to be sufficient to normalise DHEA the dose will be escalated twice to investigate whether 100 mg or 200 mg is optimal for increasing the immune response. Both 100mg and 200mg have safely been used in previous studies^{52,53}, but this has not been addressed in the context of trauma. Each group will be escalated independently of each other (**figure 2**).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Patient and public involvement (PPI)

Before the beginning of the study, patients and lay members of the 1000 elders group at the University of Birmingham were invited to group PPI sessions held by the Surgical Reconstruction and Microbiology Reconstruction Centre (SRMRC) at QEHB. During these interactive group sessions, discussions were undertaken to highlight the work that was planned to be undertaken to address previously highlighted problems in their recovery from traumatic injury. Members of the group informally looked at, developed and passed comment upon study design and patient paperwork- contemporaneous records were generated. Upon entry and active participation with the ADaPT study, patients were asked if they would like to become members of the PPI group and assist in the ongoing evaluation and future dissemination of the project. The participants will be asked to participate in a grant application should the results of this study be warranting a more extensive phase 3 multicentre trial.

Primary and secondary outcomes

The primary outcome for the study is serum DHEA after 3 days of DHEA supplementation. Previous research into DHEA levels and DHEA supplementation use DHEAS as the primary endpoint for determining whether the supplementation has been effective at raising levels. However, DHEAS levels do not act as a proxy marker for DHEA levels in the trauma population⁵¹. Utilising liquid chromatography-tandem mass spectrometry (LC-MS/MS) we have shown that DHEA and DHEAS both behave differently after trauma injury⁵⁴. Animal models of trauma have demonstrated improvements in hyperglycaemia⁵⁵, decreased mortality after trauma-induced haemorrhage⁵⁶, neurogenesis⁵⁷ and wound reperfusion⁵⁸. Human studies including a recent meta-analysis suggested that DHEA supplementation may

312 be beneficial in increasing bone mineral density (BMD)⁵⁹ in women, increasing muscle
313 strength⁶⁰, improving mood in those with moderate depression⁶¹ and adrenal insufficiency³⁸.
314 These potential restorative immunological, physiological and psychological benefits seen in
315 animals and human studies can only be investigated once the appropriate dose of DHEA to
316 restore normal levels and the most suitable administration route has been identified. We
317 know that supplementation of DHEA in healthy subjects via oral administration will result in
318 significant first-pass metabolism and, thus, more extensive conversion of DHEA to DHEAS than
319 is observed via, e.g. transdermal administration⁶² which is why we plan to compare oral vs
320 sublingual DHEA administration.

321 One potential instantaneous benefit to trauma patients, which may be observed systemically
322 after a dose of DHEA, is a positive effect upon the bactericidal function of neutrophils¹⁶ by
323 enhancing reactive oxygen species (ROS) generation via activation of NADPH oxidase⁶³.
324 Neutrophil function will therefore be investigated as a secondary outcome, with limited
325 expectations on the results of the pilot data, given that DHEA will only be administered for 3
326 days.

328 **Steroid Analysis**

329 DHEA and downstream androgens will be quantified using a validated liquid chromatography-
330 tandem mass spectrometry (LC-MS/MS) multi-steroid profiling method carried out on a
331 Waters Xevo-XS, with acquity uPLC, using a water/methanol (0.1% formic acid) gradient
332 system and a HSS T3, 1.8 µm, 1.2x50 mm column. Steroids are extracted via liquid-liquid
333 extraction using tert-butyl methyl ether (MTBE), following the addition of an internal standard
334 and protein precipitation using acetonitrile. The MTBE layer containing steroids was

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

transferred and evaporated under nitrogen then reconstituted in 125 µL of 50/50 methanol/water before analysis. Steroids will be quantified against a calibration series ranging from 0.05 to 250 ng/mL^{64–68}.

DHEAS was measured separately. 20 µL of serum was spiked with internal standard, followed by 100 µL of acetonitrile and 20 µL of ZnSO₄. The samples were then centrifuged and 100 µL of the solution was transferred to a new vial, dried and reconstituted in 200 µL of methanol/water prior to liquid chromatography tandem mass spectrometry analysis as described by Chadwick et al⁶⁹. DHEAS will be quantified against a calibration series ranging from 250 to 8000 ng/mL.

Neutrophil Function

Trauma initiates a “stress response” through the endocrine, metabolic and inflammatory systems. The primary endocrine response is to produce catecholamines and corticosteroids, raising the immune response and mobilisation of neutrophils⁷⁰. Neutrophil function will be analysed through the validated PhagoBURST™ and PhagoTest™ kits (Glycotope Biotechnology GmbH, Germany) to assess superoxide generation and phagocytosis, respectively. This gives a pilot opportunity to test whether DHEA supplementation improves the immune response and in turn, has the potential to protect against infection. However, benefits might only be detectable after a period of DHEA supplementation that is substantially longer than three days.

Pro and anti-inflammatory Cytokines

Prolonged CARS may leave the recovering patient susceptible to increased risk of late infection⁷¹. The cytokine storm of IL-6 and IL-10 have demonstrated a strong association with

the severity of injury and mortality⁷², and less so sepsis⁷³. These post-injury cytokines are also known to affect the peripheral target tissues that are involved in steroid metabolism⁷⁴. The post-injury pro and anti-inflammatory cytokines assessed have been selected based on previous work from our group⁷⁵.

Tolerance – gastric residual volume

Swallowing difficulties, facial injuries or a non-functioning gut may prohibit sublingual or oral administration and compliance to the study protocol. Therefore, an adaptable study design is needed to generate pilot data. By monitoring gastric residual volumes (GRV), a surrogate marker of gastrointestinal motility⁷⁶, we will regard a GRV persistently exceeding 250ml as intolerable.

Treatment Compliance and Evaluability

To meet study compliance and be considered evaluable, the following must be satisfied:

- Patients must be sufficiently dosed on at least one day of DHEA administration.
- If a patient fails to consume the intended IMP, or vomits within 1 hour of consuming the IMP, this dosing will be classed as *insufficient*.

Data Monitoring Committee

Data analyses will be supplied in confidence to an independent Data Monitoring Committee (DMC), which will be asked to advise on whether the accumulated data from the trial, together with the results from other relevant research, justify the continuing recruitment of further patients. The DMC will operate under a trial-specific charter based upon the template created by the Damocles Group.

Results will be provided to the DMC and discussed via teleconference at a minimum. In consultation with the trial statistician, the DMC will meet when any cohort undergoes a dose

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

escalation decision. Additional meetings may be called if recruitment is much faster than anticipated and the DMC may, at their discretion, request to meet more frequently or continue to meet following completion of recruitment. An emergency meeting may also be convened if a safety issue is identified. The DMC will report directly to the Trial Management Group. The DMC may consider recommending the discontinuation of the trial if the recruitment rate or data quality is unacceptable or if any issues are identified, which may compromise patient safety. The trial would also stop early if the interim analyses showed differences between treatments that were deemed to be convincing to the clinical community. Data monitoring members have undertaken their initial review of the first sixteen patients on the 8th November 2019. The outcome of this DMC required all-female cohorts to continue 50 mg of DHEA (both the sublingual and oral), with both male cohorts increasing to 100 mg.

Statistical Analysis

Sample Size

The maximum sample size will be 270 (six groups of 15 participants being administered three different dose-levels). However, we predict realised sample size to be smaller as there are likely to be early opportunities to escalate dose-level within a group. Following consultation with a trial statistician n = 15 was chosen to provide a modest amount of information on the primary outcome at each dose in each group while allowing recruitment to be completed in a reasonable amount of time. Statistical power calculation has not been performed as we are not applying a null hypothesis significance testing approach.

404 **Primary outcome**

405 Serum DHEA concentrations will be summarised as means and standard deviations (or
406 medians and inter-quartile ranges, if non-normal) at each time-point and dose in each cohort.
407 The observed rate of DHEA normalisation will be reported at each dose in each cohort with
408 confidence intervals calculated using Wilson's method. The cohorts sample sizes are small, so
409 these cohort-specific analyses are likely to be relatively imprecise. However, the total sample
410 size is large for the trial phase, and there is much information in the cohort structure that will
411 likely be pertinent to understanding the outcomes. Supplementary analyses to support dose
412 decisions will be provided using hierarchical regression models that analyse outcomes from
413 all cohorts and doses together while reflecting cohort memberships. These models are
414 explained below.

415

416 **Modelling serum DHEA concentrations and DHEA-normalisation probability**

417 We propose hierarchical Bayesian models to analyse serum DHEA outcomes in all cohorts
418 simultaneously. An intercept will be included to estimate the mean population-level response
419 common to all cohorts plus further terms to reflect effects for dose, patient type, and
420 administration method. Further population-level terms (also called fixed effects) will be
421 considered, including baseline DHEA level and age. Interactions will be considered, as
422 appropriate. Group-level terms (also called random effects) to reflect cohort and patient
423 heterogeneity will be considered.

424 Modestly informative or regularising priors will be used that anticipate little or no effect (i.e.
425 have a mean close to or equal to zero) but restrict the range of parameter values to those
426 that are feasible (i.e. do not place undue or unrealistic probability mass in wide distribution
427 tails). Such priors can be considered informative of scale but not location in that they do not

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

anticipate effects but rule out infeasible values. They are effective at dissuading models from over-fitting and aiding convergence in the posterior sampler when there are many parameters. Information criteria (e.g. WAIC or LOOIC) will be used to find parsimonious but informative models.

It is anticipated that: the probability of DHEA normalisation will be modelled using a generalised linear model with logit link function, and post-baseline DHEA will be modelled using a generalised linear model with identity or log (for positive data) link functions.

Interim Analysis

There will be an interim analysis presented when any cohort undergoes a dose-escalation decision, as previously described. The particular objective of this analysis will be to assess if the rate of DHEA normalisation is too low and whether there is evidence that motivates investigating a higher dose in that cohort. The primary and supporting analyses of the primary outcome will be presented, as described above.

Ethics and Dissemination

This protocol has been approved by a Research Ethics Committee (Reference 18/WM/0102) on 8th June 2018. Results will be disseminated via peer-reviewed publications and presented at national and international conferences. This will be coordinated with members of the research team, both past and present. The study investigator is responsible for communicating important protocol modifications to relevant parties.

Trial Progression

The study recruited its first patient on 2nd April 2019 and held its first data monitoring committee on 8th November 2019. Currently, there are 23 evaluable patients, with both male cohorts increasing to 100 mg. All female groups remain on 50 mg DHEA. The dose escalation also coincided with the first sponsor audit of ADaPT. Site audits will occur at times of escalation and interim analysis until the study is completed.

Figure Legends

Figure 1: A trial flowchart describing the patient journey in through the ADaPT study.

* Due to the nature of the injury, informed consent can be sought from a professional legal representative or personal legal representative if the patient does not have capacity. Consent from the patient will be obtained at the earliest opportunity by research team members.

**24hr bloods and consent will only be collected within 24hrs of injury. The omission of this sample collection does not render a patient unevaluable.

Figure 1: Indicative flowchart to explain dose escalation design of the ADaPT trial

* Cohorts: oral-male trauma, sublingual-male trauma, oral-female trauma, sublingual-female trauma, oral-female hip fracture and sublingual-female hip fracture.

Declarations

Acknowledgements: The authors thank the patients, relatives, clinical staff, research nurses and support staff in the NIHR-SRMRC at the Queen Elizabeth Hospital for their continued support for the study. NIHR-SRMRC Clinical research team members consulted at the beginning of the study are:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Gurneet Sur, Victoria Homer, Lauren Cooper; Morgan Foster, Chris McGhee, Colin Bergin, Amy Bamford, Karen Ellis, Emma Fellows, Stephanie Goundry, Elaine Spruce, Katie Moss, Tracy Mason, Christina Bratten, Liesl Despy, Samantha Harkett, Yin May, Natalie Dooley and Hazel Smith.

Competing interests: The authors have declared that no competing interests exist.

Availability of data and material: Data will be made available via online repositories.

Disclaimer: This study/project is funded by the National Institute for Health Research (NIHR) Surgical Reconstruction and Microbiology Research Centre (SRMRC). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Provenance and peer review: Not commissioned; externally peer-reviewed.

Ethics: This study has been approved by the Research Ethics Committee (REC, reference 18/WM/0102). The REC approval was gained on 8th June 2018.

Funding: For funding this research, the authors acknowledge the National Institute for Health Research Surgical Reconstruction and Microbiology Research Centre, AOUK&I Foundation and the Queen Elizabeth Hospital Charity Birmingham.

Author contributions: CB, KB and CP have prepared the manuscript. CB CP, MAF, WA, JL, CAG, AT, LC, JH, KB, AA, DB, RC, K Young, were involved in the methodological design and drafting of the trial protocol. JL, CB, CE, KM validated laboratory equipment and sample analysis for PB and PT testing. AD undertook all aspects are pharmacy procedure and IMP negotiations. K Yakoub, ET, MAF, RC, CB enrolled participants, assisted with data collection and study-

specific procedures. LC, AT, FS, WA undertook the LC-MS validation and processing of patient sex steroids samples. KB is the trial statistician who designed and wrote the analysis plan and code for randomisation of patients and times. CP, AA, DB are the trial coordinators. All authors reviewed and edited the manuscript.

Trial Sponsor: Research & Development (Governance), University Hospitals Birmingham NHS foundation Trust, Heritage Building, Queen Elizabeth Hospital, Mindlesohn Way, Edgbaston, Birmingham, B15 2GW.

References:

1. Durham RM, Moran JJ MJ. Multiple organ failure in trauma patients. [J Trauma. 2003] - PubMed result. J Trauma.
2. Lenz A, Franklin GA, Cheadle WG. Systemic inflammation after trauma. *Injury*. 2007. doi:10.1016/j.injury.2007.10.003
3. Balogh ZJ, Varga E, Tomka J, Süveges G, Tóth L, Simonka JA. The new injury severity score is a better predictor of extended hospitalization and intensive care unit admission than the injury severity score in patients with multiple orthopaedic injuries. *J Orthop Trauma*. 2003. doi:10.1097/00005131-200308000-00006
4. Foster MA, Taylor AE, Hill NE, et al. Mapping the Steroid Response to Major Trauma From Injury to Recovery: A Prospective Cohort Study. *J Clin Endocrinol Metab*. 2020;105(3). doi:10.1210/clinem/dgz302
5. Schröder J, Kahlke V, Staubach KH, Zabel P, Stüber F. Gender differences in human sepsis. *Arch Surg*. 1998. doi:10.1001/archsurg.133.11.1200

1
2
3 516 6. Frink M, Pape HC, Van Griensven M, Krettek C, Chaudry IH, Hildebrand F. Influence of
4
5
6 517 sex and age on mods and cytokines after multiple injuries. *Shock*. 2007.
7
8 518 doi:10.1097/01.shk.0000239767.64786.de
9
10
11 519 7. Hsieh YC, Frink M, Choudhry MA, Bland KI, Chaudry IH. Metabolic modulators
12
13
14 520 following trauma sepsis: Sex hormones. In: *Critical Care Medicine*. ; 2007.
15
16 521 doi:10.1097/01.CCM.0000278603.18687.4F
17
18
19 522 8. Radford DJ, Wang K, McNelis JC, et al. Dehydroepiandrosterone Sulfate Directly
20
21
22 523 Activates Protein Kinase C- β to Increase Human Neutrophil Superoxide Generation.
23
24 524 *Mol Endocrinol*. 2010. doi:10.1210/me.2009-0390
25
26
27 525 9. Duggal NA, Upton J, Phillips AC, Hampson P, Lord JM. Depressive symptoms are
28
29
30 526 associated with reduced neutrophil function in hip fracture patients. *Brain Behav*
31
32 527 *Immun*. 2013;33:173-182.
33
34
35 528 <http://linkinghub.elsevier.com/retrieve/pii/S0889159113002389>.
36
37
38 529 10. Catania RA, Angele MK, Ayala A, Cioffi WG, Bland KI, Chaudry IH.
39
40
41 530 Dehydroepiandrosterone restores immune function following trauma-haemorrhage
42
43 531 by a direct effect on T lymphocytes. *Cytokine*. 1999;11(6):443-450.
44
45 532 <http://linkinghub.elsevier.com/retrieve/pii/S1043466698904586>.
46
47
48 533 11. Eberling P, Koivisto VA. Physiological importance of dehydroepiandrosterone. *Lancet*.
49
50
51 534 1994. doi:10.1016/S0140-6736(94)92587-9
52
53
54 535 12. Hammer F, Drescher DG, Schneider SB, et al. Sex steroid metabolism in human
55
56
57 536 peripheral blood mononuclear cells changes with aging. *J Clin Endocrinol Metab*.
58
59 537 2005;90(11):6283-6289. <https://academic.oup.com/jcem/article->

- lookup/doi/10.1210/jc.2005-0915.
- 539 13. Hammer F, Subtil S, Lux P, et al. No evidence for hepatic conversion of
540 dehydroepiandrosterone (DHEA) sulfate to DHEA: in vivo and in vitro studies. *J Clin*
541 *Endocrinol Metab.* 2005;90(6):3600-3605. [https://academic.oup.com/jcem/article-](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2004-2386)
542 [lookup/doi/10.1210/jc.2004-2386](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2004-2386).
- 543 14. Arlt W, Hammer F, Sanning P, et al. Dissociation of serum dehydroepiandrosterone
544 and dehydroepiandrosterone sulfate in septic shock. *J Clin Endocrinol Metab.*
545 2006;91(7):2548-2554. [https://academic.oup.com/jcem/article-](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2005-2258)
546 [lookup/doi/10.1210/jc.2005-2258](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2005-2258).
- 547 15. Hazeldine J, Arlt W, Lord JM. Dehydroepiandrosterone as a regulator of immune cell
548 function. *J Steroid Biochem Mol Biol.* 2010;120(2):127-136.
549 <http://linkinghub.elsevier.com/retrieve/pii/S0960076009003094>.
- 550 16. Radford DJ, Wang K, McNelis JC, et al. Dehydroepiandrosterone sulfate directly
551 activates protein kinase C-beta to increase human neutrophil superoxide generation.
552 *Mol Endocrinol.* 2010;24(4):813-821. [https://academic.oup.com/mend/article-](https://academic.oup.com/mend/article-lookup/doi/10.1210/me.2009-0390)
553 [lookup/doi/10.1210/me.2009-0390](https://academic.oup.com/mend/article-lookup/doi/10.1210/me.2009-0390).
- 554 17. Oberbeck R, Kobbe P. Dehydroepiandrosterone (DHEA): A Steroid with Multiple
555 Effects. Is there Any Possible Option in the Treatment of Critical illness? *Curr Med*
556 *Chem.* 2010;17(11):1039-1047.
557 [http://www.eurekaselect.com/openurl/content.php?genre=article&issn=0929-](http://www.eurekaselect.com/openurl/content.php?genre=article&issn=0929-8673&volume=17&issue=11&spage=1039)
558 [8673&volume=17&issue=11&spage=1039](http://www.eurekaselect.com/openurl/content.php?genre=article&issn=0929-8673&volume=17&issue=11&spage=1039).
- 559 18. Butcher SK, Killampalli V, Lascelles D, Wang K, Alpar EK, Lord JM. Raised

- 560 cortisol:DHEAS ratios in the elderly after injury: potential impact upon neutrophil
561 function and immunity. *Aging Cell*. 2005;4(6):319-324.
562 <http://onlinelibrary.wiley.com/doi/10.1111/j.1474-9726.2005.00178.x/full>.
- 563 19. Phillips AC, Upton J, Duggal NA, Carroll D, Lord JM. Depression following hip fracture
564 is associated with increased physical frailty in older adults: the role of the cortisol:
565 dehydroepiandrosterone sulphate ratio. *BMC Geriatr*. 2013;13(1):60.
566 <http://bmcgeriatr.biomedcentral.com/articles/10.1186/1471-2318-13-60>.
- 567 20. Coutinho AE, Chapman KE. The anti-inflammatory and immunosuppressive effects of
568 glucocorticoids, recent developments and mechanistic insights. *Mol Cell Endocrinol*.
569 2011;335(1):2-13. <https://linkinghub.elsevier.com/retrieve/pii/S0303720710002108>.
- 570 21. Hampson P, Foster M, Taylor A, et al. The immune-endocrine mechanisms of trauma-
571 induced sarcopenia. *Endocr Abstr*. March 2014. [http://www.endocrine-](http://www.endocrine-abstracts.org/ea/0034/ea0034S5.1.htm)
572 [abstracts.org/ea/0034/ea0034S5.1.htm](http://www.endocrine-abstracts.org/ea/0034/ea0034S5.1.htm).
- 573 22. Lord JM, Midwinter MJ, Chen Y-F, et al. The systemic immune response to trauma: an
574 overview of pathophysiology and treatment. *Lancet (London, England)*.
575 2014;384(9952):1455-1465.
576 <http://linkinghub.elsevier.com/retrieve/pii/S0140673614606875>.
- 577 23. Arlt W, Haas J, Callies F, et al. Biotransformation of Oral Dehydroepiandrosterone in
578 Elderly Men: Significant Increase in Circulating Estrogens. *J Clin Endocrinol Metab*.
579 1999;84(6):2170-2176. [https://academic.oup.com/jcem/article-](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem.84.6.5789)
580 [lookup/doi/10.1210/jcem.84.6.5789](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem.84.6.5789).
- 581 24. Keane K, Hinchliffe P, Namdar N, Conceicao J, Newsholme P, Yovich J. Novel

- dehydroepiandrosterone troche supplementation improves the serum androgen profile of women undergoing in vitro fertilization. *Drug Des Devel Ther.* October 2015;5510-5569. <https://www.dovepress.com/novel-dehydroepiandrosterone-troche-supplementation-improves-the-serum-peer-reviewed-article-DDDT>.
25. Narang AS, Varia S. Role of tumor vascular architecture in drug delivery. *Adv Drug Deliv Rev.* 2011;63(8):640-658. <http://linkinghub.elsevier.com/retrieve/pii/S0169409X1100069X>.
26. Wren BG, Day RO, McLachlan AJ, Williams KM. Pharmacokinetics of estradiol, progesterone, testosterone and dehydroepiandrosterone after transbuccal administration to postmenopausal women. *Climacteric.* 2003;6(2):104-111. <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12841880&retmode=ref&cmd=prlinks>.
27. Arlt W, Callies F, Koehler I, et al. Dehydroepiandrosterone Supplementation in Healthy Men with an Age-Related Decline of Dehydroepiandrosterone Secretion. *J Clin Endocrinol Metab.* 2001;86(10):4686-4692. doi:10.1210/jcem.86.10.7974
28. Bentley C, Hazeldine J, Greig C, Lord J, Foster M. Dehydroepiandrosterone: a potential therapeutic agent in the treatment and rehabilitation of the traumatically injured patient. *Burn Trauma.* 2019;7(1):26. doi:10.1186/s41038-019-0158-z
29. World Health Organization. *Injuries and Violence. The Facts 2014.*; 2014.
30. Ward NS, Casserly B, Ayala A. The Compensatory Anti-inflammatory Response Syndrome (CARS) in Critically Ill Patients. *Clin Chest Med.* 2008;29(4):617-625. <http://www.chestmed.theclinics.com/article/S027252310800083X/fulltext>.

- 604 31. Appleton RT, Kinsella J, Quasim T. The incidence of intensive care unit-acquired
605 weakness syndromes: A systematic review. *J Intensive Care Soc.* 2015;16(2):126-136.
606 <http://journals.sagepub.com/doi/10.1177/1751143714563016>.
- 607 32. Mira JC, Brakenridge SC, Moldawer LL, Moore FA. Persistent Inflammation,
608 Immunosuppression and Catabolism Syndrome. *Crit Care Clin.* 2017;33(2):245-258.
609 <http://www.criticalcare.theclinics.com/article/S0749070416301130/fulltext>.
- 610 33. Royal College of Physicians. *National Hip Fracture Database Annual Report 2018*.;
611 2018.
- 612 34. Orentreich N, Brind JL, Rizer RL, Vogelmann JH. Age changes and sex differences in
613 serum dehydroepiandrosterone sulfate concentrations throughout adulthood. *J Clin*
614 *Endocrinol Metab.* 1984;59(3):551-555. [https://academic.oup.com/jcem/article-](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem-59-3-551)
615 [lookup/doi/10.1210/jcem-59-3-551](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem-59-3-551).
- 616 35. HORNSBY PJ. Biosynthesis of DHEAS by the Human Adrenal Cortex and Its
617 Age-Related Decline. *Ann N Y Acad Sci.* 1995. doi:10.1111/j.1749-
618 6632.1995.tb17370.x
- 619 36. Samaras N, Samaras D, Frangos E, Forster A, Philippe J. A Review of Age-Related
620 Dehydroepiandrosterone Decline and Its Association with Well-Known Geriatric
621 Syndromes: Is Treatment Beneficial? *Rejuvenation Res.* 2013.
622 doi:10.1089/rej.2013.1425
- 623 37. Orentreich N, Brind JL, Vogelmann JH, Andres R, Baldwin H. Long-term longitudinal
624 measurements of plasma dehydroepiandrosterone sulfate in normal men. *J Clin*
625 *Endocrinol Metab.* 1992;75(4):1002-1004. [https://academic.oup.com/jcem/article-](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem-75-4-1002)

- lookup/doi/10.1210/jcem.75.4.1400863.
38. Alkatib AA, Cosma M, Elamin MB, et al. A Systematic Review and Meta-Analysis of Randomized Placebo-Controlled Trials of DHEA Treatment Effects on Quality of Life in Women with Adrenal Insufficiency. *J Clin Endocrinol Metab.* 2009;94(10):3676-3681. <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=19773400&retmode=ref&cmd=prlinks>.
39. Genazzani AR, Inglese S, Lombardi I, et al. Long-term low-dose dehydroepiandrosterone replacement therapy in aging males with partial androgen deficiency. *Aging Male.* 2004;7(2):133-143. <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=15672938&retmode=ref&cmd=prlinks>.
40. Liu TC, Lin CH, Huang CY, Ivy JL, Kuo CH. Effect of acute DHEA administration on free testosterone in middle-aged and young men following high-intensity interval training. *Eur J Appl Physiol.* 2013. doi:10.1007/s00421-013-2607-x
41. Gupta B, Mittal P, Khuteta R, Bhargava A. A Comparative Study of CEE, Tibolone, and DHEA as Hormone Replacement Therapy for Surgical Menopause. *J Obstet Gynaecol India.* 2013;63(3):194-198. <http://link.springer.com/10.1007/s13224-012-0297-7>.
42. Forsblad-d'Elia H, Carlsten H, Labrie F, Konttinen YT, Ohlsson C. Low serum levels of sex steroids are associated with disease characteristics in primary Sjogren's syndrome; supplementation with dehydroepiandrosterone restores the concentrations. *J Clin Endocrinol Metab.* 2009;94(6):2044-2051. <https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2009-0106>.

- 648 43. Peuhkuri K, Vapaatalo H, Korpela R. Even low-grade inflammation impacts on small
649 intestinal function. *World J Gastroenterol*. 2010. doi:10.3748/wjg.v16.i9.1057
- 650 44. Guo Y, Hu B, Xie Y, et al. Regulation of drug-metabolizing enzymes by local and
651 systemic liver injuries. *Expert Opin Drug Metab Toxicol*. 2016.
652 doi:10.1517/17425255.2016.1139574
- 653 45. Narang N, Sharma J. Sublingual mucosa as a route for systemic drug delivery. *Int J*
654 *Pharm Pharm Sci*. 2011.
- 655 46. Chan AW, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: Defining standard
656 protocol items for clinical trials. *Ann Intern Med*. 2013. doi:10.7326/0003-4819-158-3-
657 201302050-00583
- 658 47. Duggal NA, Beswetherick A, Upton J, Hampson P, Phillips AC, Lord JM. Depressive
659 symptoms in hip fracture patients are associated with reduced monocyte superoxide
660 production. *Exp Gerontol*. 2014;54:27-34.
661 <http://linkinghub.elsevier.com/retrieve/pii/S0531556514000473>.
- 662 48. Duggal NA, Upton J, Phillips AC, Hampson P, Lord JM. Depressive symptoms post hip
663 fracture in older adults are associated with phenotypic and functional alterations in T
664 cells. *Immun Ageing*. 2014. doi:10.1186/s12979-014-0025-5
- 665 49. Pan X, Wu X, Kaminga AC, Wen SW, Liu A. Dehydroepiandrosterone and
666 Dehydroepiandrosterone Sulfate in Alzheimer's Disease: A Systematic Review and
667 Meta-Analysis. *Front Aging Neurosci*. 2019. doi:10.3389/fnagi.2019.00061
- 668 50. Trunkey DD, Lim RC. Analysis of 425 consecutive trauma fatalities: An autopsy study. *J*

- 669 *Am Coll Emerg Physicians*. 1974. doi:10.1016/S0361-1124(74)80005-5
- 670 51. Foster MA, Taylor AE, Hill NE, et al. The Endocrine and Metabolic Response in Male
671 Survivors of Major Trauma. *bioRxiv*. March 2019:577502.
672 <http://biorxiv.org/content/early/2019/03/14/577502.abstract>.
- 673 52. Morales AJ, Haubrich RH, Hwang JY, Asakura H, Yen SSC. The effect of six months
674 treatment with a 100 mg daily dose of dehydroepiandrosterone (DHEA) on circulating
675 sex steroids, body composition and muscle strength in age-advanced men and
676 women. *Clin Endocrinol (Oxf)*. 1998;49(4):421-432.
677 <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=9876338>
678 &retmode=ref&cmd=prlinks.
- 679 53. Young J, Couzinet B, Nahoul K, et al. Panhypopituitarism as a Model to Study the
680 Metabolism of Dehydroepiandrosterone (DHEA) in Humans1. *J Clin Endocrinol Metab*.
681 1997;82(8):2578-2585. doi:10.1210/jcem.82.8.4157
- 682 54. Foster M, Taylor A, Hill N, et al. The endocrine response to severe trauma: the
683 Steroids and Immunity from injury to Rehabilitation (SIR) study. *Endocr Abstr*. March
684 2014. <http://www.endocrine-abstracts.org/ea/0034/ea0034P367.htm>.
- 685 55. Sato K, Iemitsu M, Aizawa K, Mesaki N, Fujita S. Increased muscular
686 dehydroepiandrosterone levels are associated with improved hyperglycemia in obese
687 rats. *Am J Physiol Endocrinol Metab*. 2011;301(2):E274-E280.
688 <http://www.physiology.org/doi/10.1152/ajpendo.00564.2010>.
- 689 56. Angele MK, Catania RA, Ayala A, Cioffi WG, Bland KI, Chaudry IH.
690 Dehydroepiandrosterone: An inexpensive steroid hormone that decreases the

- 691 mortality due to sepsis following trauma-induced hemorrhage. *Arch Surg*.
692 1998;133(12):1281-1288.
693 <http://archsurg.jamanetwork.com/article.aspx?doi=10.1001/archsurg.133.12.1281>.
- 694 57. Karishma KK, Herbert J. Dehydroepiandrosterone (DHEA) stimulates neurogenesis in
695 the hippocampus of the rat, promotes survival of newly formed neurons and prevents
696 corticosterone-induced suppression. *Eur J Neurosci*. 2002;16(3):445-453.
697 <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12193187>
698 &retmode=ref&cmd=prlinks.
- 699 58. Lohman R, Yowell R, Barton S, Araneo B, Siemionow M. Dehydroepiandrosterone
700 protects muscle flap microcirculatory hemodynamics from ischemia/reperfusion
701 injury: An experimental in vivo study. *J Trauma - Inj Infect Crit Care*. 1997;42(1):74-80.
702 <http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00005>
703 373-199701000-00013.
- 704 59. Lin H, Li L, Wang Q, Wang Y, Wang J, Long X. A systematic review and meta-analysis of
705 randomized placebo-controlled trials of DHEA supplementation of bone mineral
706 density in healthy adults. *Gynecol Endocrinol*. 2019.
707 doi:10.1080/09513590.2019.1616175
- 708 60. Yen SS, Morales AJ, Khorram O. Replacement of DHEA in aging men and women.
709 Potential remedial effects. *Ann N Y Acad Sci*. 1995;774:128-142.
710 <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=8597453>
711 &retmode=ref&cmd=prlinks.
- 712 61. Schmidt PJ, Daly RC, Bloch M, et al. Dehydroepiandrosterone monotherapy in midlife-

- onset major and minor depression. *Arch Gen Psychiatry*. 2005;62(2):154-162.
<http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.62.2.154>.
62. Labrie F, Bélanger A, Bélanger P, et al. Metabolism of DHEA in postmenopausal women following percutaneous administration. *J Steroid Biochem Mol Biol*. 2007;103(2):178-188.
<http://linkinghub.elsevier.com/retrieve/pii/S0960076006002809>.
63. Butcher SK, Killampalli V, Lascelles D, Wang K, Alpar EK, Lord JM. Raised cortisol : DHEAS ratios in the elderly after injury: potential impact upon neutrophil function and immunity. *Aging Cell*. 2005;4(6):319-324. <http://doi.wiley.com/10.1111/j.1474-9726.2005.00178.x>.
64. Haring R, Wallaschofski H, Teumer A, et al. A SULT2A1 genetic variant identified by GWAS as associated with low serum DHEAS does not impact on the actual DHEA/DHEAS ratio. *J Mol Endocrinol*. 2013;50(1):73-77. <http://jme.endocrinology-journals.org/cgi/doi/10.1530/JME-12-0185>.
65. Büttler RM, Martens F, Fanelli F, et al. Comparison of 7 Published LC-MS/MS Methods for the Simultaneous Measurement of Testosterone, Androstenedione, and Dehydroepiandrosterone in Serum. *Clin Chem*. 2020;61(12):1475-1483.
doi:10.1373/clinchem.2015.242859
66. O'Reilly MW, Westgate CSJ, Hornby C, et al. A unique androgen excess signature in idiopathic intracranial hypertension is linked to cerebrospinal fluid dynamics. *JCI Insight*. 2019. doi:10.1172/jci.insight.125348
67. Crowley RK, Woods CP, Hughes BA, et al. Increased central adiposity and decreased

- subcutaneous adipose tissue 11 β -hydroxysteroid dehydrogenase type 1 are associated with deterioration in glucose tolerance—A longitudinal cohort study. *Clin Endocrinol (Oxf)*. 2019. doi:10.1111/cen.13939
68. O'Reilly MW, Taylor AE, Crabtree NJ, et al. Hyperandrogenemia predicts metabolic phenotype in polycystic ovary syndrome: The utility of serum androstenedione. *J Clin Endocrinol Metab*. 2014. doi:10.1210/jc.2013-3399
69. Chadwick CA, Owen LJ, Keevil BG. Development of a method for the measurement of dehydroepiandrosterone sulphate by liquid chromatography-tandem mass spectrometry. *Ann Clin Biochem*. 2005;42(Pt 6):468-474. <http://acb.sagepub.com/lookup/doi/10.1258/000456305774538175>.
70. Desborough JP. The stress response to trauma and surgery. *Br J Anaesth*. 2000;85(1):109-117. <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=10927999&retmode=ref&cmd=prlinks>.
71. Brøchner AC, Toft P. Pathophysiology of the systemic inflammatory response after major accidental trauma. *Scand J Trauma Resusc Emerg Med*. 2009;17(1):43. <http://sjtrem.biomedcentral.com/articles/10.1186/1757-7241-17-43>.
72. Stensballe J, CHRISTIANSEN M, Tønnesen E, ESPERSEN K, LIPPERT FK, RASMUSSEN LS. The early IL-6 and IL-10 response in trauma is correlated with injury severity and mortality. *Acta Anaesthesiol Scand*. 2009;53(4):515-521. <http://onlinelibrary.wiley.com/doi/10.1111/j.1399-6576.2008.01801.x/full>.
73. Frencken JF, van Vught LA, Peelen LM, et al. An Unbalanced Inflammatory Cytokine

- 757 Response Is Not Associated With Mortality Following Sepsis: A Prospective Cohort
758 Study. *Crit Care Med*. 2017;45(5):e493-e499.
759 https://journals.lww.com/ccmjournal/Fulltext/2017/05000/An_Unbalanced_Inflamm
760 [atory_Cytokine_Response_Is.36.aspx](https://journals.lww.com/ccmjournal/Fulltext/2017/05000/An_Unbalanced_Inflamm).
- 761 74. Mueller JW, Gilligan LC, Idkowiak J, Arlt W, Foster PA. The Regulation of Steroid
762 Action by Sulfation and Desulfation. *Endocr Rev*. 2015;36(5):526-563.
763 <https://academic.oup.com/edrv/article-lookup/doi/10.1210/er.2015-1036>.
- 764 75. Hazeldine J, Naumann DN, Toman E, et al. Prehospital immune responses and
765 development of multiple organ dysfunction syndrome following traumatic injury: A
766 prospective cohort study. Schreiber M, ed. *PLOS Med*. 2017;14(7):e1002338-29.
767 <http://dx.plos.org/10.1371/journal.pmed.1002338>.
- 768 76. Elke G, Felbinger TW, Heyland DK. Gastric residual volume in critically ill patients: a
769 dead marker or still alive? *Nutr Clin Pract*. 2015;30(1):59-71.
770 <http://doi.wiley.com/10.1177/0884533614562841>.




Figure 1: A trial flowchart describing the patient journey in through the ADaPT study.

* Due to the nature of the injury, informed consent can be sought from a professional legal representative or personal legal representative if the patient does not have capacity. Consent from the patient will be obtained at the earliest opportunity by research team members.

**24hr bloods and consent will only be collected within 24hrs of injury. The omission of this sample collection does not render a patient unevaluable.




Figure 2: Indicative flowchart to explain dose escalation design of the ADaPT trial * Cohorts: oral-male trauma, sublingual-male trauma, oral-female trauma, sublingual-female trauma, oral-female hip fracture and sublingual-female hip fracture.

249x287mm (300 x 300 DPI)

Reporting checklist for protocol of a clinical trial.

Based on the SPIRIT guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the SPIRITreporting guidelines, and cite them as:

Chan A-W, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, Hróbjartsson A, Mann H, Dickersin K, Berlin J, Doré C, Parulekar W, Summerskill W, Groves T, Schulz K, Sox H, Rockhold FW, Rennie D, Moher D. SPIRIT 2013 Statement: Defining standard protocol items for clinical trials. Ann Intern Med. 2013;158(3):200-207

Reporting Item			Page Number
Administrative information			
Title	#1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	1
Trial registration	#2a	Trial identifier and registry name. If not yet registered, name of intended registry	1
Trial registration: data set	#2b	All items from the World Health Organization Trial Registration Data Set	4 - ISRCTN
Protocol version	#3	Date and version identifier	10
Funding	#4	Sources and types of financial, material, and other support	23
Roles and responsibilities: contributorship	#5a	Names, affiliations, and roles of protocol contributors	23, 24

1	Roles and responsibilities:	#5b	Name and contact information for the trial sponsor	24
2	sponsor contact			
3	information			
4				
5				
6				
7				
8	Roles and responsibilities:	#5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and	23
9	sponsor and funder		interpretation of data; writing of the report; and the	
10			decision to submit the report for publication, including	
11			whether they will have ultimate authority over any of	
12			these activities	
13				
14				
15				
16				
17	Roles and responsibilities:	#5d	Composition, roles, and responsibilities of the	19
18	committees		coordinating centre, steering committee, endpoint	
19			adjudication committee, data management team, and	
20			other individuals or groups overseeing the trial, if	
21			applicable (see Item 21a for data monitoring committee)	
22				
23				
24				
25				
26	Introduction			
27				
28	Background and rationale	#6a	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention	5
29				
30				
31				
32				
33				
34				
35	Background and rationale: choice of comparators	#6b	Explanation for choice of comparators	8
36				
37				
38				
39				
40	Objectives	#7	Specific objectives or hypotheses	9
41				
42	Trial design	#8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, non-inferiority, exploratory)	9,10
43				
44				
45				
46				
47				
48				
49	Methods:			
50	Participants,			
51	interventions, and			
52	outcomes			
53				
54				
55				
56	Study setting	#9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will	7, 8, 9. The study is single site
57				
58				
59				
60				

1			be collected. Reference to where list of study sites can be	
2			obtained	
3				
4	Eligibility criteria	#10	Inclusion and exclusion criteria for participants. If	11
5			applicable, eligibility criteria for study centres and	
6			individuals who will perform the interventions (eg,	
7			surgeons, psychotherapists)	
8				
9				
10				
11	Interventions:	#11a	Interventions for each group with sufficient detail to	8, 13, Fig 2
12	description		allow replication, including how and when they will be	
13			administered	
14				
15				
16	Interventions:	#11b	Criteria for discontinuing or modifying allocated	8, 13, 19
17	modifications		interventions for a given trial participant (eg, drug dose	
18			change in response to harms, participant request, or	
19			improving / worsening disease)	
20				
21				
22				
23	Interventions:	#11c	Strategies to improve adherence to intervention	13, 19, dose-
24	adherence		protocols, and any procedures for monitoring adherence	finding and
25			(eg, drug tablet return; laboratory tests)	recorded by
26				nurses.
27				
28				
29				
30	Interventions:	#11d	Relevant concomitant care and interventions that are	Table 1
31	concomitant care		permitted or prohibited during the trial	
32				
33				
34	Outcomes	#12	Primary, secondary, and other outcomes, including the	16, 20, 21
35			specific measurement variable (eg, systolic blood	
36			pressure), analysis metric (eg, change from baseline,	
37			final value, time to event), method of aggregation (eg,	
38			median, proportion), and time point for each outcome.	
39			Explanation of the clinical relevance of chosen efficacy	
40			and harm outcomes is strongly recommended	
41				
42				
43				
44				
45	Participant timeline	#13	Time schedule of enrolment, interventions (including	9, Fig 1.
46			any run-ins and washouts), assessments, and visits for	
47			participants. A schematic diagram is highly	
48			recommended (see Figure)	
49				
50				
51				
52	Sample size	#14	Estimated number of participants needed to achieve	3
53			study objectives and how it was determined, including	
54			clinical and statistical assumptions supporting any	
55			sample size calculations	
56				
57				
58				
59				
60				

Recruitment	#15	Strategies for achieving adequate participant enrolment to reach target sample size	4,9,13,14
Methods:			
Assignment of interventions (for controlled trials)			
Allocation: sequence generation	#16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	13
Allocation concealment mechanism	#16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	13
Allocation: implementation	#16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	13, Fig 1
Blinding (masking)	#17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	n/a
Blinding (masking): emergency unblinding	#17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	n/a
Methods: Data collection, management, and analysis			
Data collection plan	#18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests)	13, 17, 18

along with their reliability and validity, if known.
Reference to where data collection forms can be found,
if not in the protocol

Data collection plan: retention	#18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	n/a – evaluable after just one dose.
Data management	#19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	13, 19
Statistics: outcomes	#20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	20,21
Statistics: additional analyses	#20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	n/a – already allocated into groups.
Statistics: analysis population and missing data	#20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	21
Methods:			
Monitoring			
Data monitoring: formal committee	#21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	19
Data monitoring: interim analysis	#21b	Description of any interim analyses and stopping guidelines, including who will have access to these	9, 14, 19

interim results and make the final decision to terminate the trial

Harms	#22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	22,13
Auditing	#23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	22
Ethics and dissemination			
Research ethics approval	#24	Plans for seeking research ethics committee / institutional review board (REC / IRB) approval	4
Protocol amendments	#25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC / IRBs, trial participants, trial registries, journals, regulators)	3, 15, 22
Consent or assent	#26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	10
Consent or assent: ancillary studies	#26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	n/a – dose-finding study.
Confidentiality	#27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	13
Declaration of interests	#28	Financial and other competing interests for principal investigators for the overall trial and each study site	22
Data access	#29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	23

1	Ancillary and post	#30	Provisions, if any, for ancillary and post-trial care, and	Fig. 1. (follow-up
2	trial care		for compensation to those who suffer harm from trial	two days post last
3			participation	dose).
4				
5				
6	Dissemination policy:	#31a	Plans for investigators and sponsor to communicate trial	22
7	trial results		results to participants, healthcare professionals, the	
8			public, and other relevant groups (eg, via publication,	
9			reporting in results databases, or other data sharing	
10			arrangements), including any publication restrictions	
11				
12				
13				
14	Dissemination policy:	#31b	Authorship eligibility guidelines and any intended use of	22
15	authorship		professional writers	
16				
17				
18	Dissemination policy:	#31c	Plans, if any, for granting public access to the full	23
19	reproducible research		protocol, participant-level dataset, and statistical code	
20				
21				
22	Appendices			
23				
24	Informed consent	#32	Model consent form and other related documentation	Due to nature of
25	materials		given to participants and authorised surrogates	patient population,
26				3 ICF and 3 PIS
27				are available.
28				
29				
30				
31	Biological specimens	#33	Plans for collection, laboratory evaluation, and storage	n/a
32			of biological specimens for genetic or molecular analysis	
33			in the current trial and for future use in ancillary studies,	
34			if applicable	
35				
36				
37				

38 None The SPIRIT checklist is distributed under the terms of the Creative Commons Attribution License CC-
39 BY-ND 3.0. This checklist can be completed online using <https://www.goodreports.org/>, a tool made by the
40 EQUATOR Network in collaboration with [Penelope.ai](#)
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

BMJ Open

A prospective, phase II, single-centre, cross-sectional, randomised study investigating Dehydroepiandrosterone supplementation and its Profile in Trauma: ADaPT

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2020-040823.R1
Article Type:	Protocol
Date Submitted by the Author:	12-Jun-2021
Complete List of Authors:	<p>Bentley, Conor; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences</p> <p>Potter, Claire; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Yakoub, Kamal; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Brock, Kristian; University of Birmingham College of Medical and Dental Sciences, D3B</p> <p>Homer, Victoria; NIHR Birmingham Liver Biomedical Research Unit Clinical Trials Group (D3B team), CRUK Clinical Trials Unit, University of Birmingham, Birmingham, UK</p> <p>Toman, Emma; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction and Microbiology Research Centre</p> <p>Taylor, Angela; University of Birmingham College of Medical and Dental Sciences, Institute of Metabolism and Systems Research</p> <p>Shaheen, Fozia; University of Birmingham College of Medical and Dental Sciences, Institute of Metabolism and Systems Research</p> <p>Gilligan, Lorna; University of Birmingham College of Medical and Dental Sciences, D3B</p> <p>Athwal, Amrita; NIHR Birmingham Liver Biomedical Research Unit Clinical Trials Group (D3B team), CRUK Clinical Trials Unit, University of Birmingham, Birmingham, UK</p> <p>Barton, Darren; NIHR Birmingham Liver Biomedical Research Unit Clinical Trials Group (D3B team), CRUK Clinical Trials Unit, University of Birmingham, Birmingham, UK</p> <p>Carrera, Ronald; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Young, Katie; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>Desai, Amisha; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre</p> <p>McGee, Kirsty; University of Birmingham College of Medical and Dental Sciences, Institute of Metabolism and Systems Research</p> <p>Ermogenous, Christos; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre;</p> <p>Sur, Gurneet; NIHR Birmingham Liver Biomedical Research Unit Clinical Trials Group (D3B team), CRUK Clinical Trials Unit, University of Birmingham, Birmingham, UK</p>

	Greig, Carolyn; University of Birmingham, School of Sport, Exercise and Rehabilitation Sciences; MRC-Arthritis Research UK Centre for Musculoskeletal Ageing Research Hazeldine, Jon; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre Arlt, Wiebke; University of Birmingham College of Medical and Dental Sciences, Institute of Metabolism and Systems Research; National Institute of Health Research Birmingham Biomedical Research Unit Lord, Janet; University of Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre; National Institute of Health Research Birmingham Biomedical Research Unit Foster, Mark; Queen Elizabeth Hospital Birmingham, NIHR Surgical Reconstruction & Microbiology Research Centre; Royal Centre for Defence Medicine
Primary Subject Heading:	Diabetes and endocrinology
Secondary Subject Heading:	Immunology (including allergy), Intensive care, Geriatric medicine, Rehabilitation medicine
Keywords:	Adult intensive & critical care < INTENSIVE & CRITICAL CARE, TRAUMA MANAGEMENT, General endocrinology < DIABETES & ENDOCRINOLOGY





I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A prospective, phase II, single-centre, cross-sectional, randomised study investigating Dehydroepiandrosterone supplementation and its Profile in Trauma: ADaPT

Conor Bentley ^{1,2,3}, Claire Potter ⁴, Kamal M. Yakoub ¹, Kristian Brock ⁴, Victoria Homer ⁴, Emma Toman ¹, Angela E. Taylor ⁵, Fozia Shaheen ⁵, Lorna C. Gilligan ⁵, Amrita Athwal ⁴, Darren Barton ⁴, Ronald Carrera ¹, Katie Young ¹, Amisha Desai ¹, Kirsty McGee ^{1,3}, Christos Ermogenous ^{1,3}, Gurneet Sur ⁴, Carolyn Greig ^{2,6}, Jon Hazeldine ^{1,3}, Wiebke Arlt ^{5,6}, Janet M. Lord ^{1,3,6}, Mark A. Foster ^{1,3,7}

¹ NIHR Surgical Reconstruction and Microbiology Research Centre, University Hospital Birmingham, Queen Elizabeth Hospital Birmingham, Birmingham B15 2WB, UK.

² School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, UK.

³ MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research, Institute of Inflammation and Ageing, Birmingham University Medical School, Birmingham B15 2TT, UK.

⁴ Diagnostics, Drugs, Devices and Biomarkers team (D³B) Cancer Research Clinical Trials Unit, University of Birmingham, Birmingham, UK.

⁵ Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK.

⁶ NIHR Biomedical Research Centre, University Hospital Birmingham and University of Birmingham, Birmingham, UK.

⁷ Royal Centre for Defence Medicine, Birmingham Research Park, Birmingham B15 2SQ, UK.

23

Correspondence to: Conor Bentley, NIHR Surgical Reconstruction and Microbiology Research
Centre, University Hospital Birmingham, Birmingham B15 2TH, UK.

email: conor.bentley@nhs.net

Tel: 0121 371 4242

28

Word Count: 4146

30

31

32

33

34

35

36

37

38

39

40

41

42

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Introduction: The improvements in short-term outcome after severe trauma achieved through early resuscitation and acute care can be offset over the following weeks by an acute systemic inflammatory response with immuneparesis leading to infection, multi-organ dysfunction/failure (MOD/MOF) and death. Serum levels of the androgen precursor dehydroepiandrosterone (DHEA) and its sulphated ester DHEAS, steroids with immune-enhancing activity, are low after traumatic injury at a time when patients are catabolic and immunosuppressed. Addressing this deficit and restoring the DHEA(S) ratio to cortisol may provide a range of physiological benefits, including immune modulatory effects.

Objective: Our primary objective is to establish a dose suitable for DHEA supplementation in patients after acute trauma to raise circulating DHEA levels to at least 15 nmol/L. Secondary objectives are to assess if DHEA supplementation has any effect on neutrophil function, metabolic and cytokine profiles and which route of administration (oral vs sub-lingual) is more effective in restoring circulating levels of DHEA, DHEAS and downstream androgens.

Methods and analysis: ADaPT is a prospective, phase II, single-centre, cross-sectional, randomised study with a planned recruitment between April 2019 and July 2021 that will investigate DHEA supplementation and its effect on serum DHEA, DHEAS and downstream androgens in trauma. A maximum of 270 patients will receive sublingual or oral DHEA at 50, 100 or 200 mg daily over 3 days. Females aged ≥ 50 years with neck of femur fracture and male and female major trauma patients, aged 16-50 years with an injury severity score ≥ 16 , will be recruited.

Ethics and dissemination: This protocol was approved by the West Midlands – Coventry and Warwickshire Research Ethics Committee (Reference 18/WM/0102) on 8th June 2018. Results

will be disseminated via peer-reviewed publications and presented at national and international conferences.

Trial registration: This trial is registered with the European Medicines Agency (EudraCT: 2016-004250-15) and ISRCTN (12961998). It has also been adopted on the National Institute of Health Research portfolio (CPMS ID:38158).

Trial Progression: The study recruited its first patient on 2nd April 2019 and held its first data monitoring committee on 8th November 2019. As of May 2020 there were 23 enrolled patients, with both male cohorts increasing to 100 mg. All female groups remain on 50 mg DHEA.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

77 **Article Summary**

78 **Strengths and limitations of this study**

79 ►► Identify the dose and route of administration needed to normalise DHEA levels in a

80 cohort that are known to have low levels, both habitually and post-traumatic injury.

81 ►► The rapid turnaround time from bedside to bench and bench to interim analysis will

82 minimise inappropriate dosing and public money expenditure on ineffective doses.

83 ►► This swift analysis will limit patient exposure to insufficient dosing and unnecessary

84 specimen collection.

85 ►► A study limitation could be its single-site design; however, this will facilitate collecting

86 sensitive immunological samples.

87 **Keywords:** steroids, geriatric trauma, major trauma, endocrine, DHEA; DHEAS; immune

88 function

89

Introduction

The improvements in short-term outcome after severe trauma achieved through early resuscitation and acute care are offset over the following weeks by an acute systemic inflammatory response with immuneparesis leading to infection, multi-organ dysfunction/failure (MOF) and death^{1,2}. The combination of excessive pro- and anti-inflammatory responses impair the rehabilitation phase of trauma, including wound healing, physical and emotional recovery³. Upregulation of glucocorticoid biosynthesis promotes a catabolic state, lasting several weeks and associated with a significant reduction in muscle mass⁴.

Analysis of gender differences in trauma has shown that women, particularly those under 30 years of age, have fewer infections and a better outcome from sepsis than men^{5,6}. The protective effects of oestrogens on immune cells and organ function highlight the potential role of sex hormones in modulating trauma related immune dysfunction⁷. Cellular immunity is also influenced by hormone production, and members of our group have shown that the adrenal response to stress, specifically the ratio of cortisol to dehydroepiandrosterone sulphate (DHEAS), is linked to neutrophil bactericidal function, specifically superoxide production^{8,9}.

Dehydroepiandrosterone (DHEA) and its sulphate ester DHEAS are the most abundant steroids in the human circulation; DHEA is a precursor of sex hormones modulating several physiologic processes including metabolism, muscle protein synthesis and cardiovascular function^{10,11}. DHEA is converted to active androgens in peripheral target cells including immune cells¹² and is also converted to DHEAS by the action of DHEA sulfotransferase (SULT2A1)¹³. We have shown that DHEA sulphation is down-regulated in acute inflammation systemic inflammatory response syndrome (SIRS) and sepsis¹⁴. Hazeldine et al reviewed the

114 numerous immune effects of DHEA ¹⁵ highlighting how DHEAS, but not DHEA, enhances
115 neutrophil superoxide generation via a protein kinase C (PKC) mediated pathway, a vital
116 immune response in fighting bacterial infections¹⁶.

117 The roles of DHEA and DHEAS in severe injury are relatively unexplored. The majority of
118 studies focus upon cortisol responses¹⁷, whereas our data suggest that it is the cortisol: DHEAS
119 ratio post-trauma that has a superior prognostic ability^{18,19}. Although critical for survival,
120 prolonged hypercortisolaemia is known to be detrimental in part due to its
121 immunosuppressive effects²⁰. This intra-adrenal shift causes decreased levels of circulating
122 testosterone and oestrogen, resulting in rapid lean body mass loss²¹, in addition to increased
123 susceptibility to infection and sepsis²². Correcting the cortisol: DHEA or cortisol: DHEAS ratio
124 via the administration of DHEA has yet to be undertaken in traumatically-injured patients
125 despite DHEA and DHEAS being below the reference ranges for 6 weeks and 6 months post-
126 injury respectively⁴.

127 DHEA is subject to first-pass metabolism in healthy individuals, which in turn may lead to a
128 non-physiological metabolism after an oral dose²³. Bypassing first-pass metabolism using a
129 sublingual or buccal preparation²⁴ should improve the bioavailability of DHEA ^{25,26}. Previous
130 studies in the healthy older people have shown that supplementation with 50 mg DHEA orally
131 once daily can restore both serum DHEA and DHEAS levels to that of men and women in the
132 third decade of life^{23,27}. Moreover, the current literature suggests that DHEA supplementation
133 may not only be beneficial for immune function but extend to bone remodelling, muscle
134 composition, psychological and neurological improvements²⁸.

135 This study will seek to estimate the optimal dose and route of administration in trauma
136 patients of DHEA to increase serum levels of DHEA and DHEAS to those seen in healthy adults.

137 We also aim to find the optimal dose to enhance immune function, as demonstrated through

changes in neutrophil phagocytosis and ROS production. The pilot data generated from the study is necessary to develop the protocol for a randomised controlled trial that will determine whether DHEA supplementation may improve outcomes in injured patients.

Rationale

Justification of the patient population

Although improvements in short-term outcomes from traumatic injury via aggressive early resuscitation and acute care, over 5 million people worldwide die each year from serious injury²⁹. With improved pre-hospital medicine mitigating the immediate threats to life, it is the following weeks after traumatic injury, that has seen the dysregulated SIRS in susceptible patients. The SIRS response is accompanied by a compensatory anti-inflammatory response (CARS)³⁰ to restore homeostasis. SIRS and CARS may progress to the persistent inflammation, immunosuppression and catabolism syndrome (PICS)^{31,32}. PICS further compounds the insult from the initial injury and results in an increased risk of infection, MOF and late deaths^{1,2}. Therefore, strategies to mitigate these detrimental outcomes for patients are needed in the short, medium and long-term care of trauma patients.

The young trauma cohort will be recruited alongside a cohort of older (≥ 50 years old) female patients who have sustained a hip fracture at the neck of femur (NOF). According to the National Hip Fracture Database (2018), the NOF burden upon the UK economy is estimated to be around £1bn per annum³³. This considerable cost is set to rise as the population ages. By the age of 40, decreasing serum levels of DHEA and DHEAS are observed in both sexes^{34,35}; a phenomenon sometimes referred to as “adrenopause”. Circulating levels of DHEA and DHEAS are lower in women than in men; however, in post-menopausal women, adrenal

androgens are a source of almost all active oestrogens³⁶. To the best of our knowledge, there has been no traumatic injury or NOF interventional studies supplementing DHEA or DHEAS, despite its therapeutic potential in the immediate to longer-term care of the young and aged trauma patient. In this pilot study, we will recruit a female NOF cohort as well as a young adult trauma cohort with both male and female patients presenting at the MTC and Intensive Care Unit (ICU) at QEHB.

Choice of treatment

The DHEA doses chosen in this study (50 mg, 100 mg and 200 mg) were selected based upon *in vivo* studies that demonstrated these doses to be both safe and effective at raising the levels of DHEA to that of healthy young adult levels³⁷. DHEA is at its highest in the third decade of life. After which there is a steady decrease over the life course in both males and females. These doses have also previously been used in adrenal insufficiency³⁸, older people³⁹, and young males⁴⁰ and females²⁴, causing transient hyperandrogenism with acne being the most frequently reported side effect^{41,42}. DHEA is activated to downstream androgens by stepwise conversion catalysed by several steroidogenic enzymes. However, we do not know whether the expression and activity of these enzymes are affected by major trauma and inflammation. There is evidence that inflammation and trauma affect DHEA sulfation^{4,14} and may shift the conversion of DHEA to a higher rate of androgen activation.

Trauma and inflammation can impact adversely on gut absorption⁴³ and hepatic first-pass metabolism⁴⁴. Therefore, we have chosen to administer DHEA as a sublingual preparation. Sub-lingual administration is commonly used by nursing staff in the ICU context, especially in those patients who have contraindications to oral administration or by patients wishing to self-administer⁴⁵.

OBJECTIVES

Primary Objective

The primary objective of this study is to establish the daily dose of DHEA that restores serum DHEA levels to at least 15 nmol/L, i.e. the mid healthy adult reference range, in trauma and hip fracture patients after 3 days of supplementation.

Secondary Objective

Secondary objectives include observing the effect of single and multiple DHEA doses on circulating DHEA, DHEAS and downstream androgens. Additionally, we will investigate whether route of DHEA administration (oral vs sublingual) modulates the profile of circulating DHEA. This will be determined by assessing the efficacy of each route via statistical modelling. The immune response to the DHEA supplementation will be assessed as a marker of potential clinical relevance.

METHODS

Trial Design

ADaPT is a prospective, randomised, open-label, trial conducted in male and female adult trauma patients and older females who have suffered a fracture of the neck of the femur. This is a single-site study with patients recruited from Queen Elizabeth Hospital, Birmingham, UK (QEHB). Three doses of DHEA will be investigated in this trial: 50, 100, 200 mg, each administered once daily for 3 days via either oral or sublingual tablets. The trial has an adaptive design in order to answer both the primary and secondary objectives, with regular interim analysis to minimise the investigation of inactive doses. The trial consists of two components. The first component of the trial is to determine the dose of DHEA needed to sufficiently raise serum DHEA levels to at least 15 nmol/L after 3 days of DHEA administration.

1
2
3 210 Based on previous work, 15 nmol/L has been selected as this is the lower acceptable level of
4
5
6 211 DHEA observed in healthy young adults. Our recent analysis of the steroid response to trauma
7
8 212 has shown that DHEA levels were very low and often undetectable for several weeks after
9
10
11 213 trauma ⁴. The second component of the trial is to investigate if DHEA will enhance neutrophil
12
13 214 function. The study was approved by The Medicines and Healthcare products Regulatory
14
15 215 Agency (MHRA) for the use of DHEA as an investigational medicinal product. Ethical approval
16
17
18 216 was obtained from the West Midlands Research Ethics Committee (Reference 18/WM/0102).
19
20
21 217 The trial will be conducted in accordance with the Declaration of Helsinki (World Medical
22
23 218 Association 2008). **Figure 1** summarises the patient pathway of the trial. The protocol (version
24
25 219 5.0, 28th June 2019) has been prepared in accordance with the SPIRIT guidelines ⁴⁶.
26
27
28 220
29
30 221
31
32

33 222 **Patient selection**
34
35 223 A maximum of 270 patients will be enrolled across three patient groups (young male trauma,
36
37 224 young female trauma and female hip fracture). These trauma patient groups have been
38
39
40 225 selected due to the immuneparesis effects caused by the acute systemic inflammatory
41
42 226 response that follows severe trauma. The hip fracture group was selected as they have low
43
44
45 227 serum DHEA and DHEAS due to adrenopause, and there are several papers showing an
46
47 228 association between HPA axis activity measures and outcomes in these patients^{9,19,47,48}.
48
49
50 229 Patients admitted to QEHB will all be pre-screened and assessed for eligibility. Patients will
51
52 230 be screened between 07:00 – 20:00, 7 days a week. Potential participants will have an
53
54 231 assessment of their medical history and current trauma injury and the eligible patients will be
55
56
57 232 recruited. The eligibility criteria have been split into trauma patients and hip fracture patients
58
59 233 (**Table 1**). The study will not exclude NOF patients with dementia where supplementation is
60

234 currently being tested in the prevention and treatment of age-related cognitive impairment
 235 without deleterious effects⁴⁹.

Table 1 Patient inclusion and exclusion criteria

Trauma patients inclusion criteria	
-	Aged 16 - 50 years of age
-	Severely injured trauma patient (Injury severity score (ISS) ≥ 16 and ≤ 50)
-	Admitted to University Hospital Birmingham within 6 days of trauma
-	Anticipated to be an inpatient for the 12-day trial period
Trauma patients exclusion criteria	
-	Ages <16 or >51
-	ISS <16 or >50
-	Isolated brain injury
-	Unlikely to survive the study period
-	Known hormone sensitive malignancy
-	Known Prostatic hypertrophy (M)
-	Female patients taking HRT medication or oral contraceptives
-	Intake of any drugs that interfere with adrenal function in the last 3 months:
Increased metabolism of glucocorticoids	
•	corticosteroids
Impaired glucocorticoid action	
Peripheral glucocorticoid insensitivity	
•	Glucocorticoid receptor antagonist—mifepristone.
•	Suppression of glucocorticoid-induced gene transcription—chlorpromazine, imipramine.
Inhibition of steroidogenic enzymes involved in cortisol production	
•	Inhibition of mitochondrial (type 1) cytochrome P450 enzymes (CYP11A1, CYP11B1, CYP11B12)—ketoconazole, fluconazole, itraconazole, etomidate, metyrapone, aminoglutethimide.
•	Inhibition of 3 β -HSD2—trilostane.
-	Pre-existing liver impairment or chronic liver failure
-	Previous or current malignancy or invasive cancer diagnosed within the past 3 years except for adequately treated basal cell and squamous cell carcinoma of the skin and in situ carcinoma of the uterine cervix
-	Pregnant and/or breast-feeding females (women of childbearing potential to complete serum pregnancy test)
-	Known hypersensitivity to the active substance or excipient
-	Known thromboembolic events in the last 12 months and any predisposition to thrombosis e.g. factor V leiden
Hip fracture patients inclusion criteria	
-	Aged 50 years and older
-	Female
-	Neck of femur fracture
-	Admitted to University Hospital Birmingham within 6 days of fracture

-	Anticipated to be an inpatient for the 12 day trial period
Hip Fracture patients exclusion criteria	
-	<50 years old
-	Unlikely to survive the study period
-	Previous or known hormone sensitive malignancy
-	Intake of any drugs that interfere with adrenal function in the last 3 months:
Increased metabolism of glucocorticoids	
•	Concomitant use reduces corticosteroid concentrations
Impaired glucocorticoid action	
Peripheral glucocorticoid insensitivity	
•	Glucocorticoid receptor antagonist—mifepristone.
•	Suppression of glucocorticoid-induced gene transcription—chlorpromazine, imipramine.
Inhibition of steroidogenic enzymes involved in cortisol production	
•	Inhibition of mitochondrial (type 1) cytochrome P450 enzymes (CYP11A1, CYP11B1, CYP11B2)—ketoconazole, fluconazole, itraconazole, etomidate, metyrapone, aminoglutethimide.
•	Inhibition of 3β-HSD2—trilostane.
-	Pre-existing liver impairment or chronic liver failure
-	Previous or current malignancy or invasive cancer diagnosed within the past 3 years except for adequately treated basal cell and squamous cell carcinoma of the skin and in situ carcinoma of the uterine cervix
-	Pregnant and/or breastfeeding (women of childbearing potential to complete serum pregnancy test)
-	Known hypersensitivity to the active substance or excipient
-	Females on Hormone Replacement Therapy medication
-	Known thromboembolic events in the last 12 months and any predisposition to thrombosis e.g. factor V leiden

Randomisation

Patients who meet the eligibility criteria and provide consent are randomised to receive DHEA via either oral tablets or sublingual tablets once daily for 3 days using a 1:1 randomisation ratio. With three populations of patients (male-trauma, female-trauma and female-hip fracture) and two routes of administration, there will be 6 groups in total. Randomisation will take place using a secure web-based tool. Nursing and medical staff will use the Clinical REsearch Tool (CREST), developed at University Hospitals Birmingham Foundation Trust (UHBFT), to randomly assign patients and provide an anonymised electronic case report form,

for trial management, data collection and adverse event reporting. The prevailing dose of DHEA (50, 100 or 200 mg) for administration in a group will be adapted in response to sequential analysis of interim outcomes. Each group will have its own dose. Blinding is not possible as the difference in DHEA delivery method is evident to both the clinician and the participant. If a contraindication to oral or the sublingual route present prior to commencing the study intervention, forced randomisation will occur.

Study Intervention

The supplementation of DHEA will commence in the second week after injury, which has previously been shown to be a time when trauma patients become maximally unwell as a result of sepsis and multiple organ failure⁵⁰. Additionally, this time point has been selected to optimise patient recruitment from both the NOF cohort and trauma patients (median stay 18 days, and 13 days respectively) both DHEA and DHEAS levels post-injury⁵¹. Recruiting an in-hospital cohort provides an opportunity to monitor patients and assess the impact that this class C controlled drug has upon the serial steroid profile and immune function during a period of vulnerability over 3 days of administration.

Participant timeline

The trial intervention will occur over three days, during the first 12 days while inpatients at QEHB. Three doses of DHEA will be investigated in all patient, and treatment will occur on day 8, 9 and 10 only. All cohorts will begin the study on 50 mg, and the dose administered to the next eligible patient to be included within a cohort will be escalated when interim analysis

1
2
3 267 determines if the primary objective has been reached. At no point will any patient escalate
4
5
6 268 once they have been allocated a dose of DHEA.
7
8

9 269 **Dose escalation**

10 270 Dose escalation for the DHEA restoration part of the trial is dependent upon the serum DHEA
11
12
13 271 levels. A dose that restores serum DHEA to ≥ 15 nmol/L (referred to as 'normalise DHEA') is
14
15
16 272 sought in at least 13-out-of-15 patients or approximately 85% of patients. The decision to
17
18 273 escalate dose in a cohort will be driven in the main by the outcomes of the patients in that
19
20 274 cohort and escalation will occur independently for each cohort. However, valuable additional
21
22
23 275 information will be garnered from the outcomes of patients in other cohorts. Flexible
24
25 276 information sharing across related groups will be achieved using hierarchical regression
26
27
28 277 models. The requirement for 13-out-of-15 patients to normalise in the final effective dose is
29
30 278 based on what was judged to be clinically important, and thus would warrant investigation of
31
32 279 the chosen dose of DHEA in a confirmatory phase III trial to ascertain superiority of outcomes
33
34
35 280 compared to standard of care.
36

37 281 To minimise the investigation of inactive doses, we propose an analysis of each dose-level in
38
39
40 282 each cohort after $n = 3-5$ evaluable patients have been assessed. It will be taken as evidence
41
42 283 that the normalisation rate is too low and that the dose should be increased if: any patient
43
44 284 fails to normalise (where $n=3$); or more than one patient fails to normalise (where $n=4$ or 5).
45
46
47 285 This rule-based analysis will provide an early opportunity to escalate in search of a more
48
49
50 286 promising dose when a probabilistic analysis will likely be under-informed due to the small
51
52 287 sample size. If the requisite number normalise, recruitment will continue in that cohort at
53
54 288 that dose-level up to a maximum of $n=15$.
55

56
57 289 Once an effective dose has been established to normalise DHEA levels within a group and at
58
59 290 least $n = 15$ have completed it, the DHEA will be escalated to the next dose to satisfy the
60

second component of the trial; to determine whether further increases in DHEA supplementation will enhance immune function. The immune response component will focus on neutrophil phagocytosis and ROS production which will involve fewer research samples. N = 15 patients (within a cohort) will complete the immune response part of the trial at each subsequent dose escalation. If 50 mg is established to be sufficient to normalise DHEA the dose will be escalated twice to investigate whether 100 mg or 200 mg is optimal for increasing the immune response. Both 100mg and 200mg have safely been used in previous studies^{52,53}, but this has not been addressed in the context of trauma. Each group will be escalated independently of each other (**figure 2**).

300

301

302 **Patient and public involvement (PPI)**

303 Before the beginning of the study, patients and lay members of the 1000 elders group at the University of Birmingham were invited to group PPI sessions held by the Surgical Reconstruction and Microbiology Reconstruction Centre (SRMRC) at QEHB. During these interactive group sessions, discussions were undertaken to highlight the work that was planned to be undertaken to address previously highlighted problems in their recovery from traumatic injury. Members of the group informally looked at, developed and passed comment upon study design and patient paperwork- contemporaneous records were generated. Upon entry and active participation with the ADaPT study, patients were asked if they would like to become members of the PPI group and assist in the ongoing evaluation and future dissemination of the project. The participants will be asked to participate in a grant

1
2
3 313 application should the results of this study be warranting a more extensive phase 3
4
5
6 314 multicentre trial.
7
8

9 315 **Primary and secondary outcomes**

10 316 The primary outcome for the study is serum DHEA after 3 days of DHEA supplementation.
11
12
13 317 Previous research into DHEA levels and DHEA supplementation use DHEAS as the primary
14
15 318 endpoint for determining whether the supplementation has been effective at raising levels.
16
17
18 319 However, DHEAS levels do not act as a proxy marker for DHEA levels in the trauma
19
20 320 population⁵¹. Utilising liquid chromatography-tandem mass spectrometry (LC-MS/MS) we
21
22 321 have shown that DHEA and DHEAS both behave differently after trauma injury⁵⁴. Animal
23
24 322 models of trauma have demonstrated improvements in hyperglycaemia⁵⁵, decreased
25
26 323 mortality after trauma-induced haemorrhage⁵⁶, neurogenesis⁵⁷ and wound reperfusion⁵⁸.
27
28
29 324 Human studies including a recent meta-analysis suggested that DHEA supplementation may
30
31 325 be beneficial in increasing bone mineral density (BMD)⁵⁹ in women, increasing muscle
32
33 326 strength⁶⁰, improving mood in those with moderate depression⁶¹ and adrenal insufficiency³⁸.
34
35
36 327 These potential restorative immunological, physiological and psychological benefits seen in
37
38 328 animals and human studies can only be investigated once the appropriate dose of DHEA to
39
40 329 restore normal levels and the most suitable administration route has been identified. We
41
42 330 know that supplementation of DHEA in healthy subjects via oral administration will result in
43
44 331 significant first-pass metabolism and, thus, more extensive conversion of DHEA to DHEAS than
45
46 332 is observed via, e.g. transdermal administration⁶² which is why we plan to compare oral vs
47
48 333 sublingual DHEA administration.
49
50
51
52
53

54 334 One potential instantaneous benefit to trauma patients, which may be observed systemically
55
56 335 after a dose of DHEA, is a positive effect upon the bactericidal function of neutrophils¹⁶ by
57
58 336 enhancing reactive oxygen species (ROS) generation via activation of NADPH oxidase⁶³.
59
60

Neutrophil function will therefore be investigated as a secondary outcome, with limited expectations on the results of the pilot data, given that DHEA will only be administered for 3 days.

Steroid Analysis

DHEA and downstream androgens will be quantified using a validated liquid chromatography-tandem mass spectrometry (LC-MS/MS) multi-steroid profiling method carried out on a Waters Xevo-XS, with acquity uPLC, using a water/methanol (0.1% formic acid) gradient system and a HSS T3, 1.8 μm , 1.2x50 mm column. Steroids are extracted via liquid-liquid extraction using tert-butyl methyl ether (MTBE), following the addition of an internal standard and protein precipitation using acetonitrile. The MTBE layer containing steroids was transferred and evaporated under nitrogen then reconstituted in 125 μL of 50/50 methanol/water before analysis. Steroids will be quantified against a calibration series ranging from 0.05 to 250 ng/mL^{64–68}.

DHEAS was measured separately. 20 μL of serum was spiked with internal standard, followed by 100 μL of acetonitrile and 20 μL of ZnSO₄. The samples were then centrifuged and 100 μL of the solution was transferred to a new vial, dried and reconstituted in 200 μL of methanol/water prior to liquid chromatography tandem mass spectrometry analysis as described by Chadwick et al⁶⁹. DHEAS will be quantified against a calibration series ranging from 250 to 8000 ng/mL.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Neutrophil Function

Trauma initiates a “stress response” through the endocrine, metabolic and inflammatory systems. The primary endocrine response is to produce catecholamines and corticosteroids, raising the immune response and mobilisation of neutrophils⁷⁰. Neutrophil function will be analysed through the validated PhagoBURST™ and PhagoTest™ kits (Glycotope Biotechnology GmbH, Germany) to assess superoxide generation and phagocytosis, respectively. This gives a pilot opportunity to test whether DHEA supplementation improves the immune response and in turn, has the potential to protect against infection. However, benefits might only be detectable after a period of DHEA supplementation that is substantially longer than three days.

Pro and anti-inflammatory Cytokines

Prolonged CARS may leave the recovering patient susceptible to increased risk of late infection⁷¹. The cytokine storm of IL-6 and IL-10 have demonstrated a strong association with the severity of injury and mortality⁷², and less so sepsis⁷³. These post-injury cytokines are also known to affect the peripheral target tissues that are involved in steroid metabolism⁷⁴. The post-injury pro and anti-inflammatory cytokines assessed have been selected based on previous work from our group⁷⁵.

Tolerance – gastric residual volume

Swallowing difficulties, facial injuries or a non-functioning gut may prohibit sublingual or oral administration and compliance to the study protocol. Therefore, an adaptable study design is needed to generate pilot data. By monitoring gastric residual volumes (GRV), a surrogate marker of gastrointestinal motility⁷⁶, we will regard a GRV persistently exceeding 250ml as intolerable.

382 Treatment Compliance and Evaluability

383 To meet study compliance and be considered evaluable, the following must be satisfied:

- 384 - Patients must be sufficiently dosed on at least one day of DHEA administration.
- 385 - If a patient fails to consume the intended IMP, or vomits within 1 hour of
386 consuming the IMP, this dosing will be classed as *insufficient*.

387 Data Monitoring Committee

388 Data analyses will be supplied in confidence to an independent Data Monitoring Committee
389 (DMC), which will be asked to advise on whether the accumulated data from the trial,
390 together with the results from other relevant research, justify the continuing recruitment of
391 further patients. The DMC will operate under a trial-specific charter based upon the template
392 created by the Damocles Group.

393 Results will be provided to the DMC and discussed via teleconference at a minimum. In
394 consultation with the trial statistician, the DMC will meet when any cohort undergoes a dose
395 escalation decision. The DMC will advise on dose escalation based on the rules as described
396 previously. Additional meetings may be called if recruitment is much faster than anticipated
397 and the DMC may, at their discretion, request to meet more frequently or continue to meet
398 following completion of recruitment. An emergency meeting may also be convened if a safety
399 issue is identified. The DMC will report directly to the Trial Management Group. The DMC may
400 consider recommending the discontinuation of the trial if the recruitment rate or data quality
401 is unacceptable or if any issues are identified, which may compromise patient safety. The trial
402 would also stop early if the interim analyses showed differences between treatments that
403 were deemed to be convincing to the clinical community. Data monitoring members have
404 undertaken their initial review of the first sixteen patients on the 8th November 2019. The

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

outcome of this DMC required all-female cohorts to continue 50 mg of DHEA (both the sublingual and oral), with both male cohorts increasing to 100 mg.

Statistical Analysis

Sample Size

The maximum sample size will be 270 (six groups of 15 participants being administered three different dose-levels). However, we predict realised sample size to be smaller as there are likely to be early opportunities to escalate dose-level within a group. Following consultation with a trial statistician n = 15 was chosen to provide a modest amount of information on the primary outcome at each dose in each group while allowing recruitment to be completed in a reasonable amount of time. Statistical power calculation has not been performed as ADaPT is a dose escalation trial and we are not applying a null hypothesis significance testing approach.

Primary outcome

Serum DHEA concentrations will be summarised as means and standard deviations (or medians and inter-quartile ranges, if non-normal) at each time-point and dose in each cohort, where a cohort is the three-way combination of patient cohort, dose and administration route. The observed rate of DHEA normalisation will be reported for each cohort with confidence intervals calculated using Wilson’s method. The cohorts sample sizes are small, so these cohort-specific analyses are likely to be relatively imprecise. However, the total sample size is large for the trial phase, and there is much information in the cohort structure that will likely be pertinent to understanding the outcomes. Supplementary analyses to support dose decisions will be provided using hierarchical regression models that analyse outcomes from

all cohorts and doses together while reflecting cohort memberships. These models are explained below.

Modelling serum DHEA concentrations and DHEA-normalisation probability

We propose hierarchical Bayesian models to analyse serum DHEA outcomes in all cohorts simultaneously. An intercept will be included to estimate the mean population-level response common to all cohorts plus further terms to reflect effects for dose, patient type, and administration method. Further population-level terms (also called fixed effects) will be considered, including baseline DHEA level and age. Interactions will be considered, as appropriate. Group-level terms (also called random effects) to reflect cohort and patient heterogeneity will be considered.

Modestly informative or regularising priors will be used that anticipate little or no effect (i.e. have a mean close to or equal to zero) but restrict the range of parameter values to those that are feasible (i.e. do not place undue or unrealistic probability mass in wide distribution tails). Such priors can be considered informative of scale but not location in that they do not anticipate effects but rule out infeasible values. They are effective at dissuading models from over-fitting and aiding convergence in the posterior sampler when there are many parameters. Information criteria (e.g. WAIC or LOOIC) will be used to find parsimonious but informative models.

It is anticipated that: the probability of DHEA normalisation will be modelled using a generalised linear model with logit link function, and post-baseline DHEA will be modelled using a generalised linear model with identity or log (for positive data) link functions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Interim Analysis

There will be an interim analysis presented when any cohort undergoes a dose-escalation decision, as previously described. The particular objective of this analysis will be to assess if the rate of DHEA normalisation is too low and whether there is evidence that motivates investigating a higher dose in that cohort. The primary and supporting analyses of the primary outcome will be presented, as described above.

Ethics and Dissemination

This protocol has been approved by a Research Ethics Committee (Reference 18/WM/0102) on 8th June 2018. Results will be disseminated via peer-reviewed publications and presented at national and international conferences. This will be coordinated with members of the research team, both past and present. The study investigator is responsible for communicating important protocol modifications to relevant parties.

Trial Progression

The study recruited its first patient on 2nd April 2019 and held its first data monitoring committee on 8th November 2019. Currently, there are 23 evaluable patients, with both male cohorts increasing to 100 mg. All female groups remain on 50 mg DHEA. The dose escalation also coincided with the first sponsor audit of ADaPT. Site audits will occur at times of escalation and interim analysis until the study is completed.

Figure Legends

Figure 1: A trial flowchart describing the patient journey in through the ADaPT study.

* Due to the nature of the injury, informed consent can be sought from a professional legal representative or personal legal representative if the patient does not have capacity. Consent from the patient will be obtained at the earliest opportunity by research team members.

**24hr bloods and consent will only be collected within 24hrs of injury. The omission of this sample collection does not render a patient unevaluable.

Figure 1: Indicative flowchart to explain dose escalation design of the ADaPT trial

* Cohorts: oral-male trauma, sublingual-male trauma, oral-female trauma, sublingual-female trauma, oral-female hip fracture and sublingual-female hip fracture.

Declarations

Acknowledgements: The authors thank the patients, relatives, clinical staff, research nurses and support staff in the NIHR-SRMRC at the Queen Elizabeth Hospital for their continued support for the study. NIHR-SRMRC Clinical research team members consulted at the beginning of the study are:

Gurneet Sur, Victoria Homer, Lauren Cooper; Morgan Foster, Chris McGhee, Colin Bergin, Amy Bamford, Karen Ellis, Emma Fellows, Stephanie Goundry, Elaine Spruce, Katie Moss, Tracy Mason, Christina Bratten, Liesl Despy, Samantha Harkett, Yin May, Natalie Dooley and Hazel Smith.

Competing interests: The authors have declared that no competing interests exist.

Availability of data and material: Data may be made available via online repositories.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Disclaimer: This study/project is funded by the National Institute for Health Research (NIHR) Surgical Reconstruction and Microbiology Research Centre (SRMRC). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Provenance and peer review: Not commissioned; externally peer-reviewed.

Ethics: This study was approved by the West Midlands – Coventry and Warwickshire Research Ethics Committee (REC, reference 18/WM/0102). The REC approval was gained on 8th June 2018.

Funding: For funding this research, the authors acknowledge the National Institute for Health Research Surgical Reconstruction and Microbiology Research Centre, AOUK&I Foundation and the Queen Elizabeth Hospital Charity Birmingham.

Author contributions: CB, KB, VH and CP have prepared the manuscript. CB, CP, MAF, WA, JL, CAG, AT, LG, JH, KB, AA, DB, RC, GS and K Young, were involved in the methodological design and drafting of the trial protocol. JL, CB, CE, KM validated laboratory equipment and sample analysis for PB and PT testing. AD undertook all aspects are pharmacy procedure and IMP negotiations. K Yakoub, ET, MAF, RC, CB enrolled participants, assisted with data collection and study-specific procedures. LG, AT, FS, WA undertook the LC-MS validation and processing of patient sex steroids samples. KB and VH are the trial statistician who designed and wrote the analysis plan and code for randomisation of patients and times. CP, AA, DB, GS are the trial coordinators. All authors reviewed and edited the manuscript.

Trial Sponsor: Research & Development (Governance), University Hospitals Birmingham NHS foundation Trust, Heritage Building, Queen Elizabeth Hospital, Mindlesohn Way, Edgbaston, Birmingham, B15 2GW.

References:

1. Durham RM, Moran JJ MJ. Multiple organ failure in trauma patients. [J Trauma. 2003] - PubMed result. J Trauma.
2. Lenz A, Franklin GA, Cheadle WG. Systemic inflammation after trauma. *Injury*. 2007. doi:10.1016/j.injury.2007.10.003
3. Balogh ZJ, Varga E, Tomka J, Süveges G, Tóth L, Simonka JA. The new injury severity score is a better predictor of extended hospitalization and intensive care unit admission than the injury severity score in patients with multiple orthopaedic injuries. *J Orthop Trauma*. 2003. doi:10.1097/00005131-200308000-00006
4. Foster MA, Taylor AE, Hill NE, et al. Mapping the Steroid Response to Major Trauma From Injury to Recovery: A Prospective Cohort Study. *J Clin Endocrinol Metab*. 2020;105(3). doi:10.1210/clinem/dgz302
5. Schröder J, Kahlke V, Staubach KH, Zabel P, Stüber F. Gender differences in human sepsis. *Arch Surg*. 1998. doi:10.1001/archsurg.133.11.1200
6. Frink M, Pape HC, Van Griensven M, Krettek C, Chaudry IH, Hildebrand F. Influence of sex and age on mods and cytokines after multiple injuries. *Shock*. 2007. doi:10.1097/01.shk.0000239767.64786.de

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

7. Hsieh YC, Frink M, Choudhry MA, Bland KI, Chaudry IH. Metabolic modulators following trauma sepsis: Sex hormones. In: *Critical Care Medicine.* ; 2007. doi:10.1097/01.CCM.0000278603.18687.4F

8. Radford DJ, Wang K, McNelis JC, et al. Dehydroepiandrosterone Sulfate Directly Activates Protein Kinase C- β to Increase Human Neutrophil Superoxide Generation. *Mol Endocrinol.* 2010. doi:10.1210/me.2009-0390

9. Duggal NA, Upton J, Phillips AC, Hampson P, Lord JM. Depressive symptoms are associated with reduced neutrophil function in hip fracture patients. *Brain Behav Immun.* 2013;33:173-182. <http://linkinghub.elsevier.com/retrieve/pii/S0889159113002389>.

10. Catania RA, Angele MK, Ayala A, Cioffi WG, Bland KI, Chaudry IH. Dehydroepiandrosterone restores immune function following trauma-haemorrhage by a direct effect on T lymphocytes. *Cytokine.* 1999;11(6):443-450. <http://linkinghub.elsevier.com/retrieve/pii/S1043466698904586>.

11. Eberling P, Koivisto VA. Physiological importance of dehydroepiandrosterone. *Lancet.* 1994. doi:10.1016/S0140-6736(94)92587-9

12. Hammer F, Drescher DG, Schneider SB, et al. Sex steroid metabolism in human peripheral blood mononuclear cells changes with aging. *J Clin Endocrinol Metab.* 2005;90(11):6283-6289. <https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2005-0915>.

13. Hammer F, Subtil S, Lux P, et al. No evidence for hepatic conversion of dehydroepiandrosterone (DHEA) sulfate to DHEA: in vivo and in vitro studies. *J Clin*

- 1
2
3 557 *Endocrinol Metab.* 2005;90(6):3600-3605. <https://academic.oup.com/jcem/article->
4
5
6 558 [lookup/doi/10.1210/jc.2004-2386](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2004-2386).
7
8
9 559 14. Arlt W, Hammer F, Sanning P, et al. Dissociation of serum dehydroepiandrosterone
10
11 560 and dehydroepiandrosterone sulfate in septic shock. *J Clin Endocrinol Metab.*
12
13 561 2006;91(7):2548-2554. <https://academic.oup.com/jcem/article->
14
15 562 [lookup/doi/10.1210/jc.2005-2258](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2005-2258).
16
17
18
19 563 15. Hazeldine J, Arlt W, Lord JM. Dehydroepiandrosterone as a regulator of immune cell
20
21 564 function. *J Steroid Biochem Mol Biol.* 2010;120(2):127-136.
22
23 565 <http://linkinghub.elsevier.com/retrieve/pii/S0960076009003094>.
24
25
26
27 566 16. Radford DJ, Wang K, McNelis JC, et al. Dehydroepiandrosterone sulfate directly
28
29 567 activates protein kinase C-beta to increase human neutrophil superoxide generation.
30
31 568 *Mol Endocrinol.* 2010;24(4):813-821. <https://academic.oup.com/mend/article->
32
33 569 [lookup/doi/10.1210/me.2009-0390](https://academic.oup.com/mend/article-lookup/doi/10.1210/me.2009-0390).
34
35
36
37 570 17. Oberbeck R, Kobbe P. Dehydroepiandrosterone (DHEA): A Steroid with Multiple
38
39 571 Effects. Is there Any Possible Option in the Treatment of Critical illness? *Curr Med*
40
41 572 *Chem.* 2010;17(11):1039-1047.
42
43 573 <http://www.eurekaselect.com/openurl/content.php?genre=article&issn=0929->
44
45 574 [8673&volume=17&issue=11&spage=1039](http://www.eurekaselect.com/openurl/content.php?genre=article&issn=0929-8673&volume=17&issue=11&spage=1039).
46
47
48
49
50
51 575 18. Butcher SK, Killampalli V, Lascelles D, Wang K, Alpar EK, Lord JM. Raised
52
53 576 cortisol:DHEAS ratios in the elderly after injury: potential impact upon neutrophil
54
55 577 function and immunity. *Aging Cell.* 2005;4(6):319-324.
56
57 578 <http://onlinelibrary.wiley.com/doi/10.1111/j.1474-9726.2005.00178.x/full>.
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

579 19. Phillips AC, Upton J, Duggal NA, Carroll D, Lord JM. Depression following hip fracture
580 is associated with increased physical frailty in older adults: the role of the cortisol:
581 dehydroepiandrosterone sulphate ratio. *BMC Geriatr.* 2013;13(1):60.
582 <http://bmcgeriatr.biomedcentral.com/articles/10.1186/1471-2318-13-60>.

583 20. Coutinho AE, Chapman KE. The anti-inflammatory and immunosuppressive effects of
584 glucocorticoids, recent developments and mechanistic insights. *Mol Cell Endocrinol.*
585 2011;335(1):2-13. <https://linkinghub.elsevier.com/retrieve/pii/S0303720710002108>.

586 21. Hampson P, Foster M, Taylor A, et al. The immune-endocrine mechanisms of trauma-
587 induced sarcopenia. *Endocr Abstr.* March 2014. [http://www.endocrine-](http://www.endocrine-abstracts.org/ea/0034/ea0034S5.1.htm)
588 [abstracts.org/ea/0034/ea0034S5.1.htm](http://www.endocrine-abstracts.org/ea/0034/ea0034S5.1.htm).

589 22. Lord JM, Midwinter MJ, Chen Y-F, et al. The systemic immune response to trauma: an
590 overview of pathophysiology and treatment. *Lancet (London, England).*
591 2014;384(9952):1455-1465.
592 <http://linkinghub.elsevier.com/retrieve/pii/S0140673614606875>.

593 23. Arlt W, Haas J, Callies F, et al. Biotransformation of Oral Dehydroepiandrosterone in
594 Elderly Men: Significant Increase in Circulating Estrogens. *J Clin Endocrinol Metab.*
595 1999;84(6):2170-2176. [https://academic.oup.com/jcem/article-](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem.84.6.5789)
596 [lookup/doi/10.1210/jcem.84.6.5789](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem.84.6.5789).

597 24. Keane K, Hinchliffe P, Namdar N, Conceicao J, Newsholme P, Yovich J. Novel
598 dehydroepiandrosterone troche supplementation improves the serum androgen
599 profile of women undergoing in vitro fertilization. *Drug Des Devel Ther.* October
600 2015:5510-5569. <https://www.dovepress.com/novel-dehydroepiandrosterone->

- troche-supplementation-improves-the-serum-peer-reviewed-article-DDDT.
25. Narang AS, Varia S. Role of tumor vascular architecture in drug delivery. *Adv Drug Deliv Rev.* 2011;63(8):640-658.
<http://linkinghub.elsevier.com/retrieve/pii/S0169409X1100069X>.
26. Wren BG, Day RO, McLachlan AJ, Williams KM. Pharmacokinetics of estradiol, progesterone, testosterone and dehydroepiandrosterone after transbuccal administration to postmenopausal women. *Climacteric.* 2003;6(2):104-111.
<http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12841880&retmode=ref&cmd=prlinks>.
27. Arlt W, Callies F, Koehler I, et al. Dehydroepiandrosterone Supplementation in Healthy Men with an Age-Related Decline of Dehydroepiandrosterone Secretion. *J Clin Endocrinol Metab.* 2001;86(10):4686-4692. doi:10.1210/jcem.86.10.7974
28. Bentley C, Hazeldine J, Greig C, Lord J, Foster M. Dehydroepiandrosterone: a potential therapeutic agent in the treatment and rehabilitation of the traumatically injured patient. *Burn Trauma.* 2019;7(1):26. doi:10.1186/s41038-019-0158-z
29. World Health Organization. *Injuries and Violence. The Facts 2014.*; 2014.
30. Ward NS, Casserly B, Ayala A. The Compensatory Anti-inflammatory Response Syndrome (CARS) in Critically Ill Patients. *Clin Chest Med.* 2008;29(4):617-625.
<http://www.chestmed.theclinics.com/article/S027252310800083X/fulltext>.
31. Appleton RT, Kinsella J, Quasim T. The incidence of intensive care unit-acquired weakness syndromes: A systematic review. *J Intensive Care Soc.* 2015;16(2):126-136.

- 622 <http://journals.sagepub.com/doi/10.1177/1751143714563016>.
- 623 32. Mira JC, Brakenridge SC, Moldawer LL, Moore FA. Persistent Inflammation,
624 Immunosuppression and Catabolism Syndrome. *Crit Care Clin*. 2017;33(2):245-258.
625 <http://www.criticalcare.theclinics.com/article/S0749070416301130/fulltext>.
- 626 33. Royal College of Physicians. *National Hip Fracture Database Annual Report 2018*.;
627 2018.
- 628 34. Orentreich N, Brind JL, Rizer RL, Vogelmann JH. Age changes and sex differences in
629 serum dehydroepiandrosterone sulfate concentrations throughout adulthood. *J Clin*
630 *Endocrinol Metab*. 1984;59(3):551-555. [https://academic.oup.com/jcem/article-](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem-59-3-551)
631 [lookup/doi/10.1210/jcem-59-3-551](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem-59-3-551).
- 632 35. HORNSBY PJ. Biosynthesis of DHEAS by the Human Adrenal Cortex and Its
633 Age-Related Decline. *Ann N Y Acad Sci*. 1995. doi:10.1111/j.1749-
634 6632.1995.tb17370.x
- 635 36. Samaras N, Samaras D, Frangos E, Forster A, Philippe J. A Review of Age-Related
636 Dehydroepiandrosterone Decline and Its Association with Well-Known Geriatric
637 Syndromes: Is Treatment Beneficial? *Rejuvenation Res*. 2013.
638 doi:10.1089/rej.2013.1425
- 639 37. Orentreich N, Brind JL, Vogelmann JH, Andres R, Baldwin H. Long-term longitudinal
640 measurements of plasma dehydroepiandrosterone sulfate in normal men. *J Clin*
641 *Endocrinol Metab*. 1992;75(4):1002-1004. [https://academic.oup.com/jcem/article-](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem.75.4.1400863)
642 [lookup/doi/10.1210/jcem.75.4.1400863](https://academic.oup.com/jcem/article-lookup/doi/10.1210/jcem.75.4.1400863).

- 643 38. Alkatib AA, Cosma M, Elamin MB, et al. A Systematic Review and Meta-Analysis of
644 Randomized Placebo-Controlled Trials of DHEA Treatment Effects on Quality of Life in
645 Women with Adrenal Insufficiency. *J Clin Endocrinol Metab.* 2009;94(10):3676-3681.
646 <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=19773400>
647 &retmode=ref&cmd=prlinks.
- 648 39. Genazzani AR, Inglese S, Lombardi I, et al. Long-term low-dose
649 dehydroepiandrosterone replacement therapy in aging males with partial androgen
650 deficiency. *Aging Male.* 2004;7(2):133-143.
651 <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=15672938>
652 &retmode=ref&cmd=prlinks.
- 653 40. Liu TC, Lin CH, Huang CY, Ivy JL, Kuo CH. Effect of acute DHEA administration on free
654 testosterone in middle-aged and young men following high-intensity interval training.
655 *Eur J Appl Physiol.* 2013. doi:10.1007/s00421-013-2607-x
- 656 41. Gupta B, Mittal P, Khuteta R, Bhargava A. A Comparative Study of CEE, Tibolone, and
657 DHEA as Hormone Replacement Therapy for Surgical Menopause. *J Obstet Gynaecol*
658 *India.* 2013;63(3):194-198. <http://link.springer.com/10.1007/s13224-012-0297-7>.
- 659 42. Forsblad-d'Elia H, Carlsten H, Labrie F, Konttinen YT, Ohlsson C. Low serum levels of
660 sex steroids are associated with disease characteristics in primary Sjogren's
661 syndrome; supplementation with dehydroepiandrosterone restores the
662 concentrations. *J Clin Endocrinol Metab.* 2009;94(6):2044-2051.
663 <https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2009-0106>.
- 664 43. Peuhkuri K, Vapaatalo H, Korpela R. Even low-grade inflammation impacts on small

- 665 intestinal function. *World J Gastroenterol*. 2010. doi:10.3748/wjg.v16.i9.1057
- 666 44. Guo Y, Hu B, Xie Y, et al. Regulation of drug-metabolizing enzymes by local and
667 systemic liver injuries. *Expert Opin Drug Metab Toxicol*. 2016.
668 doi:10.1517/17425255.2016.1139574
- 669 45. Narang N, Sharma J. Sublingual mucosa as a route for systemic drug delivery. *Int J*
670 *Pharm Pharm Sci*. 2011.
- 671 46. Chan AW, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: Defining standard
672 protocol items for clinical trials. *Ann Intern Med*. 2013. doi:10.7326/0003-4819-158-3-
673 201302050-00583
- 674 47. Duggal NA, Beswetherick A, Upton J, Hampson P, Phillips AC, Lord JM. Depressive
675 symptoms in hip fracture patients are associated with reduced monocyte superoxide
676 production. *Exp Gerontol*. 2014;54:27-34.
677 <http://linkinghub.elsevier.com/retrieve/pii/S0531556514000473>.
- 678 48. Duggal NA, Upton J, Phillips AC, Hampson P, Lord JM. Depressive symptoms post hip
679 fracture in older adults are associated with phenotypic and functional alterations in T
680 cells. *Immun Ageing*. 2014. doi:10.1186/s12979-014-0025-5
- 681 49. Pan X, Wu X, Kaminga AC, Wen SW, Liu A. Dehydroepiandrosterone and
682 Dehydroepiandrosterone Sulfate in Alzheimer's Disease: A Systematic Review and
683 Meta-Analysis. *Front Aging Neurosci*. 2019. doi:10.3389/fnagi.2019.00061
- 684 50. Trunkey DD, Lim RC. Analysis of 425 consecutive trauma fatalities: An autopsy study. *J*
685 *Am Coll Emerg Physicians*. 1974. doi:10.1016/S0361-1124(74)80005-5

- 686 51. Foster MA, Taylor AE, Hill NE, et al. The Endocrine and Metabolic Response in Male
687 Survivors of Major Trauma. *bioRxiv*. March 2019:577502.
688 <http://biorxiv.org/content/early/2019/03/14/577502.abstract>.
- 689 52. Morales AJ, Haubrich RH, Hwang JY, Asakura H, Yen SSC. The effect of six months
690 treatment with a 100 mg daily dose of dehydroepiandrosterone (DHEA) on circulating
691 sex steroids, body composition and muscle strength in age-advanced men and
692 women. *Clin Endocrinol (Oxf)*. 1998;49(4):421-432.
693 <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=9876338>
694 &retmode=ref&cmd=prlinks.
- 695 53. Young J, Couzinet B, Nahoul K, et al. Panhypopituitarism as a Model to Study the
696 Metabolism of Dehydroepiandrosterone (DHEA) in Humans1. *J Clin Endocrinol Metab*.
697 1997;82(8):2578-2585. doi:10.1210/jcem.82.8.4157
- 698 54. Foster M, Taylor A, Hill N, et al. The endocrine response to severe trauma: the
699 Steroids and Immunity from injury to Rehabilitation (SIR) study. *Endocr Abstr*. March
700 2014. <http://www.endocrine-abstracts.org/ea/0034/ea0034P367.htm>.
- 701 55. Sato K, Iemitsu M, Aizawa K, Mesaki N, Fujita S. Increased muscular
702 dehydroepiandrosterone levels are associated with improved hyperglycemia in obese
703 rats. *Am J Physiol Endocrinol Metab*. 2011;301(2):E274-E280.
704 <http://www.physiology.org/doi/10.1152/ajpendo.00564.2010>.
- 705 56. Angele MK, Catania RA, Ayala A, Cioffi WG, Bland KI, Chaudry IH.
706 Dehydroepiandrosterone: An inexpensive steroid hormone that decreases the
707 mortality due to sepsis following trauma-induced hemorrhage. *Arch Surg*.

- 1998;133(12):1281-1288.
- 709 <http://archsurg.jamanetwork.com/article.aspx?doi=10.1001/archsurg.133.12.1281>.
- 710 57. Karishma KK, Herbert J. Dehydroepiandrosterone (DHEA) stimulates neurogenesis in
- 711 the hippocampus of the rat, promotes survival of newly formed neurons and prevents
- 712 corticosterone-induced suppression. *Eur J Neurosci*. 2002;16(3):445-453.
- 713 <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12193187>
- 714 &retmode=ref&cmd=prlinks.
- 715 58. Lohman R, Yowell R, Barton S, Araneo B, Siemionow M. Dehydroepiandrosterone
- 716 protects muscle flap microcirculatory hemodynamics from ischemia/reperfusion
- 717 injury: An experimental in vivo study. *J Trauma - Inj Infect Crit Care*. 1997;42(1):74-80.
- 718 <http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00005>
- 719 373-199701000-00013.
- 720 59. Lin H, Li L, Wang Q, Wang Y, Wang J, Long X. A systematic review and meta-analysis of
- 721 randomized placebo-controlled trials of DHEA supplementation of bone mineral
- 722 density in healthy adults. *Gynecol Endocrinol*. 2019.
- 723 doi:10.1080/09513590.2019.1616175
- 724 60. Yen SS, Morales AJ, Khorram O. Replacement of DHEA in aging men and women.
- 725 Potential remedial effects. *Ann N Y Acad Sci*. 1995;774:128-142.
- 726 <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=8597453>
- 727 &retmode=ref&cmd=prlinks.
- 728 61. Schmidt PJ, Daly RC, Bloch M, et al. Dehydroepiandrosterone monotherapy in midlife-
- 729 onset major and minor depression. *Arch Gen Psychiatry*. 2005;62(2):154-162.

- 730 <http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.62.2.154>.
- 731 62. Labrie F, Bélanger A, Bélanger P, et al. Metabolism of DHEA in postmenopausal
732 women following percutaneous administration. *J Steroid Biochem Mol Biol*.
733 2007;103(2):178-188.
734 <http://linkinghub.elsevier.com/retrieve/pii/S0960076006002809>.
- 735 63. Butcher SK, Killampalli V, Lascelles D, Wang K, Alpar EK, Lord JM. Raised cortisol :
736 DHEAS ratios in the elderly after injury: potential impact upon neutrophil function
737 and immunity. *Aging Cell*. 2005;4(6):319-324. [http://doi.wiley.com/10.1111/j.1474-](http://doi.wiley.com/10.1111/j.1474-9726.2005.00178.x)
738 9726.2005.00178.x.
- 739 64. Haring R, Wallaschofski H, Teumer A, et al. A SULT2A1 genetic variant identified by
740 GWAS as associated with low serum DHEAS does not impact on the actual
741 DHEA/DHEAS ratio. *J Mol Endocrinol*. 2013;50(1):73-77. [http://jme.endocrinology-](http://jme.endocrinology-journals.org/cgi/doi/10.1530/JME-12-0185)
742 journals.org/cgi/doi/10.1530/JME-12-0185.
- 743 65. Büttler RM, Martens F, Fanelli F, et al. Comparison of 7 Published LC-MS/MS Methods
744 for the Simultaneous Measurement of Testosterone, Androstenedione, and
745 Dehydroepiandrosterone in Serum. *Clin Chem*. 2020;61(12):1475-1483.
746 doi:10.1373/clinchem.2015.242859
- 747 66. O'Reilly MW, Westgate CSJ, Hornby C, et al. A unique androgen excess signature in
748 idiopathic intracranial hypertension is linked to cerebrospinal fluid dynamics. *JCI*
749 *Insight*. 2019. doi:10.1172/jci.insight.125348
- 750 67. Crowley RK, Woods CP, Hughes BA, et al. Increased central adiposity and decreased
751 subcutaneous adipose tissue 11 β -hydroxysteroid dehydrogenase type 1 are

- associated with deterioration in glucose tolerance—A longitudinal cohort study. *Clin Endocrinol (Oxf)*. 2019. doi:10.1111/cen.13939
68. O'Reilly MW, Taylor AE, Crabtree NJ, et al. Hyperandrogenemia predicts metabolic phenotype in polycystic ovary syndrome: The utility of serum androstenedione. *J Clin Endocrinol Metab*. 2014. doi:10.1210/jc.2013-3399
69. Chadwick CA, Owen LJ, Keevil BG. Development of a method for the measurement of dehydroepiandrosterone sulphate by liquid chromatography-tandem mass spectrometry. *Ann Clin Biochem*. 2005;42(Pt 6):468-474.
<http://acb.sagepub.com/lookup/doi/10.1258/000456305774538175>.
70. Desborough JP. The stress response to trauma and surgery. *Br J Anaesth*. 2000;85(1):109-117.
<http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=10927999&retmode=ref&cmd=prlinks>.
71. Brøchner AC, Toft P. Pathophysiology of the systemic inflammatory response after major accidental trauma. *Scand J Trauma Resusc Emerg Med*. 2009;17(1):43.
<http://sjtrem.biomedcentral.com/articles/10.1186/1757-7241-17-43>.
72. Stensballe J, CHRISTIANSEN M, Tønnesen E, ESPERSEN K, LIPPERT FK, RASMUSSEN LS. The early IL-6 and IL-10 response in trauma is correlated with injury severity and mortality. *Acta Anaesthesiol Scand*. 2009;53(4):515-521.
<http://onlinelibrary.wiley.com/doi/10.1111/j.1399-6576.2008.01801.x/full>.
73. Frencken JF, van Vught LA, Peelen LM, et al. An Unbalanced Inflammatory Cytokine Response Is Not Associated With Mortality Following Sepsis: A Prospective Cohort

- 774 Study. *Crit Care Med*. 2017;45(5):e493-e499.
- 775 https://journals.lww.com/ccmjournal/Fulltext/2017/05000/An_Unbalanced_Inflamm
- 776 [atory_Cytokine_Response_Is.36.aspx](https://journals.lww.com/ccmjournal/Fulltext/2017/05000/An_Unbalanced_Inflamm).
- 777 74. Mueller JW, Gilligan LC, Idkowiak J, Arlt W, Foster PA. The Regulation of Steroid
- 778 Action by Sulfation and Desulfation. *Endocr Rev*. 2015;36(5):526-563.
- 779 <https://academic.oup.com/edrv/article-lookup/doi/10.1210/er.2015-1036>.
- 780 75. Hazeldine J, Naumann DN, Toman E, et al. Prehospital immune responses and
- 781 development of multiple organ dysfunction syndrome following traumatic injury: A
- 782 prospective cohort study. Schreiber M, ed. *PLOS Med*. 2017;14(7):e1002338-29.
- 783 <http://dx.plos.org/10.1371/journal.pmed.1002338>.
- 784 76. Elke G, Felbinger TW, Heyland DK. Gastric residual volume in critically ill patients: a
- 785 dead marker or still alive? *Nutr Clin Pract*. 2015;30(1):59-71.
- 786 <http://doi.wiley.com/10.1177/0884533614562841>.




Figure 1: A trial flowchart describing the patient journey in through the ADaPT study.

* Due to the nature of the injury, informed consent can be sought from a professional legal representative or personal legal representative if the patient does not have capacity. Consent from the patient will be obtained at the earliest opportunity by research team members.

**24hr bloods and consent will only be collected within 24hrs of injury. The omission of this sample collection does not render a patient unevaluable.




Figure 2: Indicative flowchart to explain dose escalation design of the ADaPT trial * Cohorts: oral-male trauma, sublingual-male trauma, oral-female trauma, sublingual-female trauma, oral-female hip fracture and sublingual-female hip fracture.

249x287mm (600 x 600 DPI)

Reporting checklist for protocol of a clinical trial.

Based on the SPIRIT guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the SPIRITreporting guidelines, and cite them as:

Chan A-W, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, Hróbjartsson A, Mann H, Dickersin K, Berlin J, Doré C, Parulekar W, Summerskill W, Groves T, Schulz K, Sox H, Rockhold FW, Rennie D, Moher D. SPIRIT 2013 Statement: Defining standard protocol items for clinical trials. Ann Intern Med. 2013;158(3):200-207

Reporting Item			Page Number
Administrative information			
Title	#1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	1
Trial registration	#2a	Trial identifier and registry name. If not yet registered, name of intended registry	1
Trial registration: data set	#2b	All items from the World Health Organization Trial Registration Data Set	4 - ISRCTN
Protocol version	#3	Date and version identifier	10
Funding	#4	Sources and types of financial, material, and other support	23
Roles and responsibilities: contributorship	#5a	Names, affiliations, and roles of protocol contributors	23, 24

1	Roles and responsibilities:	#5b	Name and contact information for the trial sponsor	24
2	sponsor contact			
3	information			
4				
5				
6				
7				
8	Roles and responsibilities:	#5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	23
9	sponsor and funder			
10				
11				
12				
13				
14				
15				
16				
17	Roles and responsibilities:	#5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	19
18	committees			
19				
20				
21				
22				
23				
24				
25				
26	Introduction			
27				
28	Background and rationale	#6a	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention	5
29				
30				
31				
32				
33				
34				
35	Background and rationale: choice of comparators	#6b	Explanation for choice of comparators	8
36				
37				
38				
39				
40	Objectives	#7	Specific objectives or hypotheses	9
41				
42	Trial design	#8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, non-inferiority, exploratory)	9,10
43				
44				
45				
46				
47				
48				
49	Methods:			
50	Participants, interventions, and outcomes			
51				
52				
53				
54				
55				
56	Study setting	#9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will	7, 8, 9. The study is single site
57				
58				
59				
60				

1			be collected. Reference to where list of study sites can be	
2			obtained	
3				
4	Eligibility criteria	#10	Inclusion and exclusion criteria for participants. If	11
5			applicable, eligibility criteria for study centres and	
6			individuals who will perform the interventions (eg,	
7			surgeons, psychotherapists)	
8				
9				
10				
11	Interventions:	#11a	Interventions for each group with sufficient detail to	8, 13, Fig 2
12	description		allow replication, including how and when they will be	
13			administered	
14				
15				
16	Interventions:	#11b	Criteria for discontinuing or modifying allocated	8, 13, 19
17	modifications		interventions for a given trial participant (eg, drug dose	
18			change in response to harms, participant request, or	
19			improving / worsening disease)	
20				
21				
22				
23	Interventions:	#11c	Strategies to improve adherence to intervention	13, 19, dose-
24	adherence		protocols, and any procedures for monitoring adherence	finding and
25			(eg, drug tablet return; laboratory tests)	recorded by
26				nurses.
27				
28				
29				
30	Interventions:	#11d	Relevant concomitant care and interventions that are	Table 1
31	concomitant care		permitted or prohibited during the trial	
32				
33				
34	Outcomes	#12	Primary, secondary, and other outcomes, including the	16, 20, 21
35			specific measurement variable (eg, systolic blood	
36			pressure), analysis metric (eg, change from baseline,	
37			final value, time to event), method of aggregation (eg,	
38			median, proportion), and time point for each outcome.	
39			Explanation of the clinical relevance of chosen efficacy	
40			and harm outcomes is strongly recommended	
41				
42				
43				
44				
45	Participant timeline	#13	Time schedule of enrolment, interventions (including	9, Fig 1.
46			any run-ins and washouts), assessments, and visits for	
47			participants. A schematic diagram is highly	
48			recommended (see Figure)	
49				
50				
51				
52	Sample size	#14	Estimated number of participants needed to achieve	3
53			study objectives and how it was determined, including	
54			clinical and statistical assumptions supporting any	
55			sample size calculations	
56				
57				
58				
59				
60				

Recruitment	#15	Strategies for achieving adequate participant enrolment to reach target sample size	4,9,13,14
Methods:			
Assignment of interventions (for controlled trials)			
Allocation: sequence generation	#16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	13
Allocation concealment mechanism	#16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	13
Allocation: implementation	#16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	13, Fig 1
Blinding (masking)	#17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	n/a
Blinding (masking): emergency unblinding	#17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	n/a
Methods: Data collection, management, and analysis			
Data collection plan	#18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests)	13, 17, 18

along with their reliability and validity, if known.
Reference to where data collection forms can be found,
if not in the protocol

Data collection plan: retention	#18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	n/a – evaluable after just one dose.
Data management	#19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	13, 19
Statistics: outcomes	#20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	20,21
Statistics: additional analyses	#20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	n/a – already allocated into groups.
Statistics: analysis population and missing data	#20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	21
Methods:			
Monitoring			
Data monitoring: formal committee	#21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	19
Data monitoring: interim analysis	#21b	Description of any interim analyses and stopping guidelines, including who will have access to these	9, 14, 19

interim results and make the final decision to terminate the trial

Harms	#22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	22,13
Auditing	#23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	22
Ethics and dissemination			
Research ethics approval	#24	Plans for seeking research ethics committee / institutional review board (REC / IRB) approval	4
Protocol amendments	#25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC / IRBs, trial participants, trial registries, journals, regulators)	3, 15, 22
Consent or assent	#26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	10
Consent or assent: ancillary studies	#26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	n/a – dose-finding study.
Confidentiality	#27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	13
Declaration of interests	#28	Financial and other competing interests for principal investigators for the overall trial and each study site	22
Data access	#29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	23

1	Ancillary and post	#30	Provisions, if any, for ancillary and post-trial care, and	Fig. 1. (follow-up
2	trial care		for compensation to those who suffer harm from trial	two days post last
3			participation	dose).
4				
5				
6	Dissemination policy:	#31a	Plans for investigators and sponsor to communicate trial	22
7	trial results		results to participants, healthcare professionals, the	
8			public, and other relevant groups (eg, via publication,	
9			reporting in results databases, or other data sharing	
10			arrangements), including any publication restrictions	
11				
12				
13				
14	Dissemination policy:	#31b	Authorship eligibility guidelines and any intended use of	22
15	authorship		professional writers	
16				
17				
18	Dissemination policy:	#31c	Plans, if any, for granting public access to the full	23
19	reproducible research		protocol, participant-level dataset, and statistical code	
20				
21				
22	Appendices			
23				
24	Informed consent	#32	Model consent form and other related documentation	Due to nature of
25	materials		given to participants and authorised surrogates	patient population,
26				3 ICF and 3 PIS
27				are available.
28				
29				
30				
31	Biological specimens	#33	Plans for collection, laboratory evaluation, and storage	n/a
32			of biological specimens for genetic or molecular analysis	
33			in the current trial and for future use in ancillary studies,	
34			if applicable	
35				
36				
37				

38 None The SPIRIT checklist is distributed under the terms of the Creative Commons Attribution License CC-
39 BY-ND 3.0. This checklist can be completed online using <https://www.goodreports.org/>, a tool made by the
40 EQUATOR Network in collaboration with [Penelope.ai](#)
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60